

## Therapeutic Class Overview

### Attention-Deficit/Hyperactivity Disorder (ADHD) Agents

#### INTRODUCTION

- Attention-deficit/hyperactivity disorder (ADHD) is the most common neurodevelopmental disorder among children, with an estimated prevalence of up to 10% of school-age children in the United States (U.S.). It is more common in boys than girls and frequently persists into adulthood (*Centers for Disease Control and Prevention [CDC] 2021, Feldman et al 2014*). Epidemiologic studies of adult ADHD have estimated the current prevalence to be 4.4% in the U.S. (*Bukstein 2021*).
  - In children, this chronic disorder is characterized by symptoms of hyperactivity, impulsivity, and/or inattention. These symptoms affect cognitive, academic, behavioral, emotional, and social functioning (*Krull 2019*). Common comorbid psychiatric disorders include oppositional defiant disorder, conduct disorder, depression, anxiety disorder, and learning disabilities (*Krull 2021a*). Approximately 20% of children with ADHD develop chronic tic disorders and approximately 50% of children with chronic tics or Tourette syndrome have comorbid ADHD (*Krull 2021c*).
  - ADHD in adults is characterized by symptoms of inattention, impulsivity, and restlessness. Impairment in executive function and emotional dysregulation frequently occur. Common comorbid psychiatric disorders include mood and anxiety disorders, substance use disorder, and intermittent explosive disorder (*Bukstein 2021*).
- For children < 17 years of age, the Diagnostic and Statistical Manual of Mental Disorders Fifth Edition (DSM-5) diagnosis of ADHD requires ≥ 6 symptoms of hyperactivity and impulsivity or ≥ 6 symptoms of inattention. For adolescents ≥ 17 years of age and adults, ≥ 5 symptoms of hyperactivity and impulsivity or ≥ 5 symptoms of inattention are required.
  - The symptoms of hyperactivity/impulsivity or inattention must occur often; be present in more than 1 setting; persist for at least 6 months; be present before the age of 12 years; impair function in academic, social, or occupational activities; and be excessive for the developmental level of the child.
  - Other physical, situational, or mental health conditions that could account for the symptoms must be excluded.
- Treatment of ADHD may involve behavioral/psychologic interventions, medication, and/or educational interventions, alone or in combination (*Krull 2020*).
  - For preschool children (age 4 through 5 years), behavioral therapy is considered the first-line treatment; when medication is necessary, methylphenidate is generally recommended.
  - For children and adolescents with moderate to severe ADHD, medication and behavioral therapy are recommended. In general, stimulants are the first-line agents; however, nonstimulant medications may be more appropriate for certain children.
    - About 30% of patients do not respond to or may not tolerate the initial stimulant treatment. At least one-half of children who do not respond to one type of stimulant will respond to the other. If there is still no improvement, consideration should be given to switching to or adding a nonstimulant ADHD medication (*Krull 2021b*).
- Multiple agents are currently approved by the Food and Drug Administration (FDA) for the treatment of ADHD. They include central nervous system (CNS) stimulants (amphetamine- and methylphenidate-based formulations), as well as nonstimulants: 2 selective norepinephrine reuptake inhibitors (SNRIs) atomoxetine and viloxazine extended-release (ER); and 2 alpha<sub>2</sub>-adrenergic agonists clonidine ER, guanfacine ER.
  - Due to the potential for abuse, the stimulant agents are classified as Schedule II controlled substances.
  - Several stimulants are also approved for the treatment of narcolepsy and exogenous obesity; the use of stimulants for the treatment of obesity will not be covered in this review. Lisdexamfetamine dimesylate is the only FDA-approved drug for the treatment of binge eating disorder (BED).
- Medispan Classes: ADHD Agents – Amphetamines, Dexmethylphenidate, Methylphenidate, Selective Alpha-Adrenergic Agonists, Selective Norepinephrine Reuptake Inhibitor

**Table 1. Medications Included Within Class Review**

<b>Drug</b>	<b>Generic Availability</b>
<b>Stimulants</b>	
Evekeo (amphetamine sulfate)	✓
Evekeo ODT (amphetamine sulfate)	-
Azstarys (serdexmethylphenidate/dexmethylphenidate)	-
Adderall (mixed amphetamine salts)	✓
Focalin (dexmethylphenidate hydrochloride [HCl])	✓
ProCentra (dextroamphetamine sulfate)	✓
Zenzedi (dextroamphetamine sulfate)	✓
Desoxyn (methamphetamine HCl)	✓
methylphenidate HCl chewable tablets	✓
Methylin Oral Solution (methylphenidate HCl)	✓
Ritalin (methylphenidate HCl)	✓
Dexedrine Spansule (dextroamphetamine sulfate sustained-release)	✓
Adzenys XR-ODT (amphetamine ER)	-
Dyanavel XR (amphetamine ER)	-
Adderall XR (mixed amphetamine salts ER)	✓
Mydayis (mixed amphetamine salts ER)	-
Focalin XR (dexmethylphenidate HCl ER)	✓
Vyvanse (lisdexamfetamine dimesylate)	-
Adhansia XR (methylphenidate HCl ER)	-
Aptensio XR (methylphenidate HCl ER)	✓
Concerta (methylphenidate HCl ER)	✓
Cotempla XR-ODT (methylphenidate ER)	-
Jornay PM (methylphenidate HCl ER)	-
methylphenidate HCl ER (CD)	✓
methylphenidate HCl ER	✓
QuilliChew ER (methylphenidate HCl ER)	-
Quillivant XR (methylphenidate HCl ER)	-
Relexxii (methylphenidate HCl ER) (72 mg)	✓
Ritalin LA (methylphenidate HCl ER)	✓
Daytrana (methylphenidate transdermal system)	-
<b>Nonstimulants</b>	
Strattera (atomoxetine HCl)	✓
Kapvay (clonidine HCl ER)	✓
Intuniv (guanfacine HCl ER)	✓
Qelbree (viloxazine ER)	-

*(Drugs@FDA 2021, Orange Book: Approved Drug Products with Therapeutic Equivalence Evaluations 2021, Clinical Pharmacology 2021)*

**INDICATIONS**

**Table 2. Food and Drug Administration Approved Indications**

Indication	ADHD*	ADHD, as an integral part of a total treatment program which typically includes other remedial measures (psychological, educational, and social) for a stabilizing effect in pediatric patients with a behavioral syndrome characterized by the following group of developmentally inappropriate symptoms: moderate to severe distractibility, short attention span, hyperactivity, emotional lability, and impulsivity. The diagnosis of this syndrome should not be made with finality when these symptoms are only of comparatively recent origin. Nonlocalizing (soft) neurological signs, learning disability, and abnormal electroencephalogram (EEG) may or may not be present, and a diagnosis of CNS dysfunction may or may not be warranted.*	Treatment of ADHD as monotherapy and as adjunctive therapy to stimulant medications	Narcolepsy**	Exogenous obesity, as a short term (a few weeks) adjunct in a regimen of weight reduction based on caloric restriction for patients refractory to alternative therapy (eg, repeated diets, group programs, and other drugs).†	Moderate to severe BED in adults
Evekeo (amphetamine sulfate)		✓		✓	✓	
Evekeo ODT (amphetamine sulfate)	✓					
Adzenys XR-ODT, Dyanavel XR (amphetamine)	✓					
Adderall (mixed amphetamine salts)	✓			✓		
Adderall XR, Mydayis (mixed amphetamine salts ER)	✓					
Strattera (atomoxetine HCl)	✓					
Kapvay (clonidine HCl ER)			✓			
Focalin (dexamethylphenidate IR); Focalin XR (dexamethylphenidate ER)	✓					
ProCentra, Zenzedi (dextroamphetamine sulfate IR); Dexedrine Spansule (dextroamphetamine sulfate SR)		✓		✓		
Intuniv (guanfacine HCl ER)			✓			
Vyvanse (lisdexamfetamine dimesylate)	✓					✓

Desoxyn (methamphetamine HCl)		✓			
Ritalin (methylphenidate HCl IR); methylphenidate HCl chewable tablets		✓		✓	
Methylin Oral Solution; methylphenidate ER tablets	✓			✓	
Adhansia XR, Aptensio XR, Concerta, Cotelpla XR-ODT, Daytrana, Jornay PM, QuilliChew ER, Quillivant XR, Relexxii, Ritalin LA (methylphenidate ER)	✓				
Azstarys (serdexmethylphenidate/dexmethylphenidate)	✓				
Qelbree (viloxazine ER)	✓				

(Prescribing Information: Adderall 2020, Adderall XR 2020, Adhansia XR 2021, Adzenys XR-ODT 2018, Aptensio XR 2021, Azstarys 2021, Concerta 2021, Cotelpla XR-ODT 2021, Daytrana 2021, Desoxyn 2019, Dexedrine Spansule 2019, Dyanavel XR 2021, Evekeo 2019, Evekeo ODT 2021, Focalin 2021, Focalin XR 2021, Intuniv 2020, Jornay PM 2021, Kapvay 2020, Mydayis 2020, Methylin Oral Solution 2021, methylphenidate chewable tablets 2021, methylphenidate ER 2021, methylphenidate ER (CD) 2021, ProCentra 2017, Qelbree 2021, QuilliChew ER 2021, Quillivant XR 2021, Relexxii 2019, Ritalin 2021, Ritalin LA 2021, Strattera 2020, Vyvanse 2021, Zenzedi 2021)

\* Adderall, Evekeo, ProCentra, and Zenzedi are approved for use in children 3 years of age and older. Evekeo ODT is approved for use in patients 3 to 17 years of age. Daytrana, Desoxyn, Dexedrine Spansule, Intuniv, and Kapvay are approved for use in children 6 years of age and older. Adderall XR, Adhansia XR, Adzenys XR-ODT, Aptensio XR, Azstarys, Dyanavel XR, Focalin, Focalin XR, Jornay PM, methylphenidate ER (CD), methylphenidate ER, Methylin Oral Solution, methylphenidate chewable tablets, QuilliChew ER, Quillivant XR, Ritalin, Strattera, and Vyvanse are approved for use in patients 6 years of age and older. Cotelpla XR-ODT and Qelbree are approved for use in pediatric patients 6 to 17 years of age. Ritalin LA is approved for use in pediatric patients 6 to 12 years of age. Concerta and Relexxii are approved for use in children 6 years of age and older, adolescents, and adults up to 65 years of age. Mydayis is approved for use in patients 13 years of age and older.

\*\*These drugs are approved for use in patients 6 years of age and older.

†These drugs are not recommended for use in children under 12 years of age for treatment of exogenous obesity. The limited usefulness of these products should be weighed against possible risks inherent in use of the drugs.

- Limitation of use:
  - Aptensio XR: Pediatric patients younger than 6 years of age experienced higher plasma exposure than patients 6 years and older at the same dose and high rates of AEs, most notably weight loss.
  - Lisdexamfetamine: Pediatric patients younger than 6 years of age experienced more long-term weight loss than patients 6 years and older. Lisdexamfetamine is not indicated or recommended for weight loss. Use of other sympathomimetic drugs for weight loss has been associated with serious cardiovascular (CV) adverse events (AEs). The safety and effectiveness of this drug for the treatment of obesity have not been established.
  - Mydayis: Pediatric patients 12 years and younger experienced higher plasma exposure than patients 13 years and older at the same dose and experienced higher rates of AEs, mainly insomnia and decreased appetite.
- Information on indications, mechanism of action, pharmacokinetics, dosing, and safety has been obtained from the prescribing information for the individual products, except where noted otherwise.

## CLINICAL EFFICACY SUMMARY

- Randomized trials, systematic reviews, and meta-analyses have found stimulants, SNRIs (atomoxetine, viloxazine ER), and alpha<sub>2</sub>-adrenergic agonists (clonidine ER, guanfacine ER) to be more efficacious than placebo in reducing the core symptoms of ADHD in children and adolescents.
  - Evekeo (amphetamine sulfate) was approved based on a randomized, double-blind (DB), multicenter (MC), placebo-controlled (PC) laboratory classroom study that was conducted in 107 children between the ages of 6 and 12 years (Childress et al 2015). The study found Evekeo to be associated with significant improvements in the average

Swanson, Kotkin, Agler, M-Flynn, and Pelham (SKAMP) combined score compared to placebo (least squares [LS] mean difference -7.9; 95% CI, -10.1 to -5.6;  $p < 0.0001$ ).

- Evekeo ODT, an orally disintegrating amphetamine tablet, was approved under the 505(b)(2) regulatory pathway. The safety and effectiveness of Evekeo ODT for the treatment of ADHD was established based on an adequate and well-controlled study of Evekeo (*Childress et al 2015*).
- Cotelpla XR-ODT, a new methylphenidate ER orally disintegrating tablet formulation, was approved based on a randomized, DB, MC, PC laboratory classroom study (*Childress et al 2017*) ( $N = 87$ ) which found that the average SKAMP-combined score was significantly better for Cotelpla XR-ODT than for placebo (LS mean 14.3 [95% CI, 12.2 to 16.4] vs 25.3 [9% CI, 23.0 to 27.6], respectively;  $p < 0.0001$ ).
- Adhansia XR, a recently approved methylphenidate ER capsule, was approved via the 505(b)(2) regulatory pathway, and its efficacy was supported by 4 clinical studies in patients with ADHD including 2 studies conducted in adults, 1 study in adolescents 12 to 17 years of age, and 1 study in pediatric patients 6 to 12 years of age (*Adhansia XR FDA Clinical Review 2019*):
  - One randomized, DB, MC, PC 4-week study conducted in 368 adult patients with ADHD evaluated the safety and efficacy of 4 doses of Adhansia XR (25, 45, 70, and 100 mg) compared to placebo. The primary endpoint, change in the ADHD-Rating Scale (ADHD-RS)-5 total score from baseline to Week 5, was significantly improved compared to placebo in the Adhansia XR 45 mg group (LS mean difference, -6.9; 95% CI, -11.5 to -2.2;  $p = 0.0013$ ), 100 mg group (LS mean difference, -8.1; 95% CI, -12.9 to -3.2;  $p = 0.0002$ ), and when combining all dosage groups compared to placebo (LS mean difference, -4.7; 95% CI, -7.7 to -1.6;  $p = 0.0026$ ). No significant difference was seen in the 25 mg or 70 mg groups compared to placebo.
  - A second randomized, DB, crossover, PC study was conducted in 45 adults in an adult workplace environment (*Adhansia XR FDA Clinical Review 2019, Wigal et al 2020*). The study aimed to assess efficacy parameters for Adhansia XR vs placebo over 16 hours post-dose. Patients were titrated to an optimal dose of Adhansia XR (either 25, 35, 45, 55, 70, 85, or 100 mg) during an open-label (OL) treatment period between 2 and 7 weeks, then entered into a 1-week PC, DB treatment phase prior to the adult workplace environment session, followed by a 7-day washout period between crossover periods, then another 1-week treatment phase followed by another adult workplace environment session. The primary endpoint was the average Permanent Product Measure of Performance (PERMP) score for various time points up to 16 hours post-dose. When combining data from all time points, patients treated with Adhansia XR had significant improvements in the PERMP score compared to placebo (LS mean difference, 13.05; 95% CI, 3.88 to 22.23;  $p = 0.0064$ ).
  - A 4-week randomized, DB, PC trial assessed efficacy of Adhansia XR in 354 adolescent patients 12 to 17 years of age (*Adhansia XR FDA Clinical Review 2019*). The study compared Adhansia XR 25, 45, 70, and 85 mg to placebo and found significant improvements in the ADHD-5-RS score from baseline to Week 5 in adolescents treated with Adhansia XR 45 mg (LS mean difference, -5.4; 95% CI, -9.2 to -1.6;  $p = 0.0052$ ), 70 mg (LS mean difference, -5.2; 95% CI, -9.0 to -1.4;  $p = 0.0069$ ), and when combining all dosage groups compared to placebo (LS mean difference, -4.3; 95% CI, -7.3 to -1.3;  $p = 0.0049$ ). Adolescents treated with Adhansia XR 25 or 85 mg did not achieve significant improvements in the ADHD-5-RS score compared to placebo.
    - A fourth study, which included a 6-week OL dose optimization period (majority of patients received between 45 and 55 mg of Adhansia XR) followed by a 1-week DB, PC study, was conducted to assess the efficacy of Adhansia XR in 147 children 6 to 12 years of age in an analog classroom setting. The primary endpoint, average SKAMP-C score (taken at various time points up to 13 hours post-dose), was significantly improved in children treated with Adhansia XR compared to placebo (LS mean difference, -8.6; 95% CI, -10.6 to -6.6).
- Jornay PM, an ER methylphenidate capsule formulation, was approved based on the results of 2 clinical studies conducted in patients 6 to 12 years of age with ADHD:
  - The first study was a 6-week OL dose-optimization study, followed by a 1-week DB, PC withdrawal phase where patients were randomized to continue treatment with Jornay PM or switch to placebo (*Childress et al 2020, Jornay PM Prescribing Information 2021*). The study, which was conducted in an analog classroom setting and included 117 children aged 6 to 12 years, found that Jornay PM was associated with a significant reduction in the SKAMP symptom score over a 12-hour period (LS mean difference, -5.9; 95% CI, -9.1 to -2.7).
  - A randomized, DB, MC, PC, parallel group, forced-dose titration trial was conducted over 3 weeks in 161 children 6 to 12 years of age with ADHD (*Pliszka et al 2017*). The study found that 40 to 80 mg/day of Jornay PM achieved significant improvements vs placebo in ADHD symptoms (LS mean ADHD rating scale-IV, 24.1 vs 31.2;  $p = 0.002$ ) at 3 weeks. Significant improvements were also seen vs placebo in key secondary outcomes including at-home



early morning and late afternoon/evening functional impairment at 3 weeks. The most commonly reported treatment-emergent AEs were insomnia and decreased appetite.

- o Mydayis, a mixed amphetamine salts product, was approved for the treatment of ADHD based on the results of 5 MC, DB, PC, randomized controlled trials (RCTs): 3 in adults and 2 in pediatric patients 13 to 17 years of age. The studies found that Mydayis demonstrated a statistically significant treatment effect compared with placebo on various ADHD outcomes measures (eg, ADHD-RS score, PERMP score) (*Mydayis Prescribing Information 2020, Weisler et al 2017, Wigal et al 2018a, Wigal et al 2018b, Wigal et al 2019*) (see results below in Table 3 below). An additional 6-week, randomized, PC, DB, forced dose titration trial in 411 adults with ADHD similarly found that Mydayis significantly improved ADHD-RS-IV scores compared to placebo (LS mean treatment difference for all Mydayis doses combined vs placebo, -10.6; 95% CI, -13.2 to -8.0;  $p < 0.0001$ ) (*Frick et al 2020*).

**Table 3. Summary of Primary Efficacy Results for Mydayis**

Study Number (Age range)	Primary Endpoint	Treatment Group	Mean Baseline Score (SD)	LS Mean Change from Baseline	Placebo-subtracted Difference (95% CI)
<b>Adult Studies</b>					
Study 1 (18 to 55 years)	ADHD-RS	Mydayis 12.5 mg/day <sup>§</sup>	39.8 (6.38)	-18.5	-8.1 (-11.7 to -4.4)
		Mydayis 37.5 mg/day <sup>§</sup>	39.9 (7.07)	-23.8	
		Placebo	40.5 (6.52)	-10.4	
Study 2 (18 to 55 years)	Average PERMP	Mydayis 50 mg/day <sup>§</sup>	239.2 (75.6) <sup>†</sup>	293.23*	18.38 (11.28 to 25.47)
		Placebo	249.6 (76.7) <sup>†</sup>	274.85*	
Study 3 (18 to 55 years)	Average PERMP	Mydayis 25 mg/day <sup>§</sup>	217.5 (59.6) <sup>†</sup>	267.96*	19.29 (10.95 to 27.63)
		Placebo	226.9 (61.7) <sup>†</sup>	248.67*	
<b>Pediatric Studies</b>					
Study 4 (13 to 17 years) <sup>‡</sup>	ADHD-RS-IV	Mydayis 12.5 to 25 mg/day <sup>§</sup>	36.7 (6.15)	-20.3	-8.7 (-12.6 to -4.8)
		Placebo	38.3 (6.67)	-11.6	
Study 5 (13 to 17 years)	Average PERMP	Mydayis 25 mg/day <sup>§</sup>	214.5 (87.8) <sup>†</sup>	272.67*	41.26 (32.24 to 50.29)
		Placebo	228.7 (101) <sup>†</sup>	231.41*	

SD = standard deviation; LS = least squares; CI = confidence interval

<sup>†</sup>Pre-dose PERMP total score

\*LS mean for PERMP is post-dose average score over all sessions of the treatment day, rather than change from baseline

<sup>‡</sup>Results are for a subgroup of study 4 and not the total population

<sup>§</sup>Doses statistically significant for placebo

- o Azstarys, a combination of serdexmethylphenidate and dexamethylphenidate, was approved based on results from a randomized, DB, PC analog classroom study (*Azstarys Prescribing Information 2021*). A total of 150 patients aged 6 to 12 years were enrolled. Following an OL, 3-week dose titration phase, patients were randomly assigned during a 1-week parallel treatment period to either the optimized dose Azstarys or placebo. After 1 week, evaluations were done using the SKAMP rating scale over 13 hours in a classroom setting. Mean change in SKAMP from baseline (primary outcome) was significantly greater with Azstarys compared with placebo (placebo-subtracted difference -5.4; 95% CI, -7.1 to -3.7). The efficacy of Azstarys in adults and pediatric patients 13 to 17 years of age was established by pharmacokinetic bridging between Azstarys and Focalin XR (dexamethylphenidate ER) capsules.
- o Qelbree (viloxazine ER), an SNRI, was shown to be superior to placebo in 3 DB, MC, randomized, PC trials in patients with ADHD.
  - Trial 1 enrolled 313 patients aged 6 to 11 years who were randomized to treatment with viloxazine ER 200 or 400 mg or placebo once daily for 8 weeks (*Nasser 2021b*). Improvements in ADHD-RS-5 total scores were reported, with LS mean changes from baseline of -17.6, -17.5 and -11.7 for viloxazine ER 200 mg, 400 mg, and placebo, respectively ( $p < 0.05$  for both comparisons to placebo).
  - Trial 2 enrolled 477 patients aged 6 to 11 years who were randomized to either viloxazine ER 100 mg or 200 mg or placebo once daily for 6 weeks (*Nasser 2020*). LS mean changes from baseline in ADHD-RS-5 total scores were

- 16.6, -17.7, and -10.9 for viloxazine ER 100 mg, 200 mg, and placebo, respectively ( $p < 0.05$  and  $p < 0.0001$  for viloxazine ER 100 mg and 200 mg vs placebo, respectively).
- A third trial evaluated viloxazine ER in 310 patients aged 12 to 17 years of age who were randomized to viloxazine ER 200 mg, 400 mg, or placebo (Nasser 2021a). After 6 weeks of treatment, viloxazine ER 200 mg and 400 mg resulted in LS mean changes from baseline in ADHD-RS-5 total scores of -16.0, -16.5, and -11.4 for viloxazine ER 200 mg, 400 mg, and placebo, respectively ( $p < 0.05$  vs placebo for both comparisons).
  - A systematic (Cochrane) review of 185 RCTs (Storebø et al 2015) (N = 12,245) in children and adolescents with ADHD found that methylphenidate may improve teacher-rated ADHD symptoms, teacher-reported general behavior, and parent-reported quality of life (QOL) vs placebo. However, the evidence was of low quality.
  - An RCT called the Preschool ADHD Treatment Study (PATS) (Greenhill et al 2006) evaluated the efficacy of methylphenidate immediate-release (IR) in 303 preschool children with ADHD and found that it demonstrated significant reductions on ADHD symptom scales; however, the effect sizes (0.4 to 0.8) were smaller than those generally reported for school-age children.
  - A systematic (Cochrane) review of 23 PC, RCTs (Punja et al 2016) (N = 2675) found that amphetamines were effective at improving the core symptoms of ADHD, but they were also associated with a higher risk of AEs compared to placebo. There was no evidence that one kind of amphetamine was better than another and there was no difference between short-acting and long-acting formulations.
  - A meta-analysis of 25 DB, PC, RCTs (Schwartz et al 2014) (N = 3928) in children and adolescents with ADHD found atomoxetine to be superior to placebo for overall ADHD symptoms, with a medium effect size (-0.64).
  - A meta-analysis of 25 RCTs (all rated as low or very low quality evidence) in children with autism and concurrent ADHD symptoms concluded that methylphenidate and atomoxetine both reduced parent-rated hyperactivity and inattention (Rodrigues et al 2021). Methylphenidate also reduced teacher-rated hyperactivity and inattention, but atomoxetine only reduced teacher-rated inattention.
  - A meta-analysis of 12 RCTs (Hirota et al 2014) (N = 2276) in pediatric patients with ADHD found that alpha<sub>2</sub>-adrenergic agonists were significantly superior to placebo for overall ADHD symptoms both as monotherapy and, to a lesser extent, as augmentation therapy to stimulants.
    - Meta-analytic results failed to demonstrate a significant difference in efficacy between alpha<sub>2</sub>-adrenergic agonists. In sub-analyses of individual formulations, the ER formulations separated robustly from placebo whereas the IR formulations did not separate from placebo.
  - A systematic review of 16 RCTs and 1 meta-analysis (Chan et al 2016) (N = 2668) found evidence supporting the use of methylphenidate ER and amphetamine ER formulations, atomoxetine, and guanfacine ER for the treatment of ADHD in adolescents. For the primary outcome measure of mean change in ADHD-RS total symptom score, both stimulant and nonstimulant medications led to clinically significant reductions of 14.93 to 24.60 points.
  - For the treatment of ADHD in children and adolescents, stimulants typically have a slightly larger treatment effect size (standardized mean difference [SMD]) than nonstimulants (approximately 1.0 vs approximately 0.7 for both atomoxetine and alpha<sub>2</sub>-adrenergic agonists). However, there is insufficient evidence to definitively conclude that one stimulant is more efficacious than another (Krull 2021b, (Wolraich et al 2019).
    - An Agency for Healthcare Research and Quality (AHRQ) review of 78 studies (Jadad et al 1999) evaluating the efficacy of various interventions for the treatment of ADHD in children and adults found few, if any, differences between methylphenidate and dextroamphetamine.
    - A meta-analysis of 23 DB, PC trials (Faraone 2010a) comparing the efficacy of methylphenidate and amphetamine formulations found that amphetamine products may be moderately more efficacious than methylphenidate products.
    - A DB, PC, RCT (Newcorn et al 2008) (N = 516) comparing the efficacy of atomoxetine vs methylphenidate ER (osmotic-release formulation) in patients 6 to 16 years of age with ADHD found that both drugs were superior to placebo in terms of response rate, and that methylphenidate ER was superior to atomoxetine.
    - A meta-analysis of 29 DB, PC trials (Faraone et al 2006) evaluated the efficacy of various medications (methylphenidate and amphetamine compounds, atomoxetine, pemoline [no longer available in the U.S.], bupropion, and modafinil) for the treatment of ADHD. The effect sizes for nonstimulant medications were significantly less than those for IR stimulants or long-acting stimulants. The 2 classes of stimulant medications did not differ significantly from one another.
    - A meta-analysis of 28 DB, PC, RCTs (Stuhec et al 2015) (N = 4699) compared the efficacy of various medications for the treatment of ADHD in children and adolescents. Efficacy in reducing ADHD symptoms compared to placebo was small for bupropion (SMD, -0.32; 95% CI, -0.69 to 0.05), modest for atomoxetine (SMD, -0.68; 95% CI, -0.76 to -0.59)

- and methylphenidate (SMD, -0.75; 95% CI, -0.98 to -0.52), and highest for lisdexamfetamine (SMD, -1.28; 95% CI, -1.84 to -0.71).
- A network meta-analysis and mixed treatment comparison of 36 RCTs (*Joseph et al 2017*) evaluating the comparative efficacy and safety of ADHD pharmacotherapies in children and adolescents found that lisdexamfetamine had greater efficacy than guanfacine ER, atomoxetine, and methylphenidate ER. Guanfacine ER had a high posterior probability of being more efficacious than atomoxetine, but their credible intervals overlapped.
  - A network meta-analysis of 48 DB, RCTs (*Padilha et al 2018*) compared the safety and efficacy of various ADHD medications in children and adolescents. Of the 12 trials that were evaluated for efficacy, analysis was performed using the Clinical Global Impression Improvement (CGI-I) scale for 3 drugs, which showed that methylphenidate was more effective than atomoxetine (MD, 3.15; 95% CI, 0.75 to 13.71) and guanfacine (MD, 1.92; 95% CI, 0.64 to 5.94). Thirty-three trials were evaluated for safety. Ranking of AEs showed that lisdexamfetamine was more likely to cause sleep disorders, loss of appetite, and behavior problems compared to other treatments.
  - Alpha<sub>2</sub>-adrenergic agonists have been associated with improvements in ADHD symptoms and comorbid tics.
    - A meta-analysis of 9 DB, PC, RCTs (*Bloch et al 2009*) (N = 477) was conducted to determine the relative efficacy of different medications in treating ADHD and tic symptoms in children with both Tourette syndrome and ADHD.
    - Methylphenidate seemed to offer the greatest improvement of ADHD symptoms and did not seem to worsen tic symptoms.
    - Alpha<sub>2</sub>-adrenergic agonists offered the best combined improvement in both tic and ADHD symptoms.
    - Atomoxetine significantly improved both tic and ADHD severity compared to placebo.
    - One small study found that tic severity was significantly increased with higher doses of dextroamphetamine treatment.
    - A Cochrane review of 8 RCTs (*Osland et al 2018*) including 510 children with both ADHD and a chronic tic disorder found low-quality evidence for improvement of ADHD symptoms with methylphenidate, atomoxetine, and clonidine, and very low-quality evidence for desipramine, dextroamphetamine, guanfacine, and deprenyl. Tic symptoms improved with guanfacine, desipramine, methylphenidate, clonidine, and a combination of methylphenidate and clonidine. The authors noted that in 1 study with a short duration (3 weeks), high doses of dextroamphetamine worsened tics.
  - There are limited efficacy data regarding the treatment of ADHD in the adult population. Comparison of effect sizes in clinical trials suggests that stimulant medications are more efficacious in adult ADHD than nonstimulants.
    - In a meta-analysis of 12 clinical trials (*Cunill et al 2013*) (N = 3375) comparing atomoxetine with placebo in adult ADHD, atomoxetine led to a modestly greater reduction in ADHD symptom severity, but was associated with higher all-cause discontinuation.
    - A meta-analysis (*Faraone 2010b*) of 19 randomized trials of 13 medications for adult ADHD found a greater average effect size for reduction in ADHD symptoms in patients receiving short- and long-acting stimulant medications (vs placebo; 0.86 and 0.73, respectively) compared with patients receiving nonstimulant medication (vs placebo; 0.39). No difference in effect size was found between short- and long-acting stimulants.
    - A meta-analysis of 20 randomized trials (*Stuhec et al 2019*) compared the efficacy, acceptability, and tolerability of lisdexamfetamine, mixed amphetamine salts, methylphenidate, and modafinil in the treatment of ADHD in adults. The highest effect size in reducing ADHD symptoms was found with lisdexamfetamine (SMD -0.89; 95% CI, -1.09 to -0.70), while moderate reductions in symptoms were seen with mixed amphetamine salts (SMD -0.64; 95% CI, -0.83 to -0.45) and methylphenidate (SMD -0.50; 95% CI, -0.58 to -0.41). No efficacy was reported with modafinil.
    - A Cochrane review of 19 studies (*Castells et al 2018*, N = 2521) comparing dextroamphetamine, lisdexamfetamine, and mixed amphetamine salts for the treatment of ADHD in adults found that overall, amphetamines reduced the patient- and clinician-rated severity of ADHD symptoms compared to placebo; however, they did not improve retention in treatment. Amphetamines were associated with an increased proportion of patients who withdrew because of AEs. When comparing different types of amphetamines, lisdexamfetamine and mixed amphetamine salts reduced the severity of ADHD symptoms as rated by clinicians, but dextroamphetamine did not. No differences in any outcome were found when comparing immediate- and sustained-release formulations.
    - A systematic review and network meta-analysis (*Elliot et al 2020*) of 81 RCTs compared methylphenidate, atomoxetine, dexamfetamine, lisdexamfetamine, guanfacine, mixed amphetamine salts, modafinil, and bupropion for the treatment of ADHD in adults. Treatment with any ADHD pharmacotherapy was associated with statistically significant improvement in patient-reported clinical response vs placebo. When drugs were analyzed individually, only atomoxetine was found to significantly improve patient-reported clinical response compared to placebo (mean difference [MD], -5.9; 95% CI, -12.6 to -0.4). Atomoxetine (MD, -3.7; 95% CI, -6.7 to -0.9), sustained-release methylphenidate (MD, -5.7; 95% CI, -11.2 to -0.3), and low-dose methylphenidate (MD, -10.4; 95% CI, -19.0 to -2.1)



were found to improve clinician-assessed clinical response compared to placebo. No significant differences were observed between individual medications when response was considered as a continuous outcome.

- Another meta-analysis (*Cortese et al 2018*) of 133 RCTs comparing the use of amphetamines, atomoxetine, bupropion, clonidine, guanfacine, methylphenidate, and modafinil for the treatment of ADHD found that all drugs were superior to placebo for ADHD core symptoms as rated by clinicians in children and adolescents, and all drugs except for modafinil were more efficacious than placebo in adults.
  - When comparing the various drugs based on teachers' ratings in children and adolescents, only methylphenidate and modafinil were found to be more efficacious than placebo.
  - In head-to-head comparisons, differences in efficacy based on clinicians' ratings were found, favoring amphetamines over modafinil (SMD, -0.39; 95% CI -0.67 to -0.12), atomoxetine (SMD, -0.46; 95% CI, -0.65 to -0.27), and methylphenidate (SMD, -0.24; 95% CI, -0.44 to -0.05) in children and adolescents. Efficacy results based on clinicians' ratings were similar for adults, and favored amphetamines over modafinil (SMD, -0.94; 95% CI -1.43 to -0.46), atomoxetine (SMD, -0.34; 95% CI, -0.58 to -0.10), and methylphenidate (SMD, -0.29; 95% CI, -0.54 to -0.05).
- Lisdexamfetamine dimesylate has demonstrated efficacy in the treatment of BED. Direct comparison trials between lisdexamfetamine and other drugs used off-label to treat BED are lacking.
  - In 2 Phase 3, 12-week, randomized, DB, PC trials (*McElroy et al 2016*) (N = 773) in patients with moderate to severe BED, lisdexamfetamine-treated patients had a statistically significantly greater reduction from baseline in mean number of binge days per week at week 12 vs placebo (treatment difference in study 1: -1.35; 95% confidence interval [CI], -1.70 to -1.01; study 2: -1.66; 95% CI, -2.04 to -1.28; both p < 0.001).
    - A 12-month, OL extension study (*Gasior et al 2017*) (N = 599) in adults with BED found that the long-term safety and tolerability of lisdexamfetamine were generally consistent with the safety profile observed in 3 previous short-term trials in BED as well as its established profile for ADHD. Common treatment-emergent AEs included dry mouth, headache, insomnia, and upper respiratory tract infection. Weight loss and increases in blood pressure and pulse rate were also observed.
  - In a phase 3, DB, randomized, PC, withdrawal study (*Hudson et al 2017*) (N = 418) in adults with moderate to severe BED, responders to lisdexamfetamine during a 12-week OL phase were randomized to placebo or continued lisdexamfetamine during a 26-week, DB phase. The percentage of patients meeting relapse criteria was 3.7% with lisdexamfetamine vs 32.1% with placebo; time to relapse statistically favored lisdexamfetamine (p < 0.001). The hazard ratio (HR) was 0.09 (95% CI, 0.04 to 0.23).
  - A systematic review and meta-analysis of 14 clinical and 7 preclinical trials concluded that lisdexamfetamine effectively treats BED and reduces both symptoms (MD, 0.93; 95% CI, 0.74 to 1.12) and body weight (based on systematic review only) (*Schneider et al 2021*).
  - A systematic review and meta-analysis of 9 waitlist-controlled psychological trials and 25 PC trials evaluating pharmacologic (n = 19) or combination (n = 6) treatment for BED (*Brownley et al 2016*) found that therapist-led cognitive behavioral therapy (CBT), lisdexamfetamine, and second-generation antidepressants (SGAs) increased binge-eating abstinence (relative risk [RR], 4.95 [95% CI, 3.06 to 8.00], 2.61 [95% CI, 2.04 to 3.33], and 1.67 [95% CI, 1.24 to 2.26], respectively), while lisdexamfetamine and SGAs decreased binge-eating frequency (MD in days/week, -1.35 [95% CI, -1.77 to -0.93] and -0.67 [95% CI, -1.26 to -0.09], respectively). Topiramate and other forms of CBT also increased abstinence and reduced binge-eating frequency.
  - A 2018 systematic review and meta-analysis of 45 RCTs (*Ghaderi et al 2018*) compared various psychological, pharmacological, and combined treatments for BED, and found moderate support for the efficacy of CBT and CBT-guided self-help (moderate quality of evidence), and low-quality evidence to support interpersonal psychotherapy, selective serotonin reuptake inhibitors (SSRIs), and lisdexamfetamine for the cessation of or reduction in the frequency of binge eating. Only lisdexamfetamine showed a modest effect on weight loss (SMD for body mass index -5.23; 95% CI, -6.52 to -3.94).

## CLINICAL GUIDELINES

### ADHD

- Several clinical guidelines have provided recommendations on the treatment of ADHD in children and adolescents.
  - According to the American Academy of Pediatrics (AAP) guidelines (*Wolraich et al 2019*), the evidence is particularly strong for stimulant medications, and sufficient but less strong for atomoxetine, guanfacine ER, and clonidine ER (in that order; newer agents such as serdexmethylphenidate/dexmethylphenidate [Azstarys] and viloxazine [Qelbree] are

not addressed in the current guidelines). Guanfacine ER and clonidine ER have evidence to support their use as adjunctive therapy with stimulant medications. Methylphenidate is recommended for preschool-aged children who have had an inadequate response to behavioral interventions.

- The Society for Developmental and Behavioral Pediatrics guideline on assessment and treatment of children and adolescents with complex ADHD states that treatment should aim to improve functional impairment and include skill development in self-management strategies (*Barbarese et al 2020*). Multimodal treatment with both behavioral and pharmacologic therapies may be needed. Specific pharmacologic classes are discussed in the context of learning disorder, for which the guideline recommends both stimulants and atomoxetine, with stimulants having a greater strength of evidence, and autism, for which a stimulant is recommended first followed by an alpha<sub>2</sub>-adrenergic agonist or atomoxetine. Stimulant use is also endorsed in children with intellectual disability, tics, anxiety or depression, and disruptive behavior disorders.
- The Medical Letter recommends that treatment of ADHD in school-age children or adults should begin with a stimulant, either a methylphenidate- or amphetamine-based formulation (*Med Lett Drugs Ther 2020*). Mixing short- and long-acting stimulants can be helpful to achieve an immediate effect for early-morning school classes or for reducing rebound irritability or overactivity, especially in the evening. Nonstimulants can be used in combination with stimulants or when stimulants are contraindicated, ineffective, or not tolerated.
- According to the American Academy of Neurology guidelines for treatment of tics (*Pringsheim et al 2019*), physicians should counsel individuals with tics and comorbid ADHD that alpha<sub>2</sub>-adrenergic agonists may provide benefit for both conditions. Alpha<sub>2</sub>-adrenergic agonists and topiramate should be prescribed for the treatment of tics when the benefits of treatment outweigh the risks, while antipsychotics and botulinum toxin may be prescribed when the benefits outweigh the risks.
- The American Academy of Child and Adolescent Psychiatry (AACAP) practice parameter for the treatment of children and adolescents with tic disorders (*Murphy et al 2013*) states that alpha<sub>2</sub>-adrenergic agonists have demonstrated an effect size of 0.5 for the amelioration of tics and may be preferred by some prescribers over antipsychotics due to their relatively favorable AE profile.

### Narcolepsy

- The American Academy of Sleep Medicine (AASM) practice parameters (*Maski et al 2021*) recommend various drugs for the treatment of daytime sleepiness in adults due to narcolepsy including modafinil, pitolisant, sodium oxybate, solriamfetol (strongly recommended), and armodafinil, dextroamphetamine, and methylphenidate (conditionally recommended). Idiopathic hypersomnia in adults should be treated with modafinil (strongly recommended), clarithromycin, methylphenidate, pitolisant, or sodium oxybate (conditionally recommended). Recommended therapies for children with narcolepsy include modafinil and sodium oxybate (both conditionally recommended).

### BED

- According to the American Psychiatric Association (APA) practice guidelines on eating disorders (*Yager et al 2006, Yager et al 2012* [guideline watch update], now categorized as a legacy guideline), treatment of BED may include the following:
  - Nutritional rehabilitation and counseling
  - Psychosocial treatment
    - CBT, behavior therapy, dialectical behavior therapy (DBT), and interpersonal therapy (IPT) have all been associated with binge frequency reduction rates of 67% or more and significant abstinence rates during active treatment.
    - Self-help programs using self-guided, professionally designed manuals have been effective in reducing the symptoms of BED in the short-run for some patients and may have long-term benefit.
  - Medications
    - Antidepressant treatment is associated with short-term reductions in binge-eating but generally does not result in substantial weight loss. SSRIs have the fewest difficulties with AEs and the most evidence for efficacy when used at the high end of the recommended dose range.
    - Topiramate can reduce bingeing and decrease weight, but its use may be limited by AEs.
  - Combination psychotherapy and pharmacotherapy
    - For most patients, adding antidepressant therapy to a behavioral weight control and/or CBT regimen does not have a significant effect on binge suppression.
    - Although limited evidence is available, combined treatment is frequently used in clinical practice.

- The American Association of Clinical Endocrinologists and the American College of Endocrinology (AACE/ACE) guidelines for medical care of patients with obesity (*Garvey et al 2016*) recommend the following for patients with overweight or obesity who have BED:
  - Patients should be treated with a structured behavioral/lifestyle program, combined with CBT or other psychological interventions
  - Treatment with orlistat or approved medications containing topiramate or bupropion may be considered in conjunction with structured lifestyle therapy, CBT, and/or psychological interventions
- The Task Force on Eating Disorders of the World Federation of Societies of Biological Psychiatry (*Aigner et al 2011*) concluded that for the treatment of BED, grade A evidence supports the use of imipramine (moderate risk-benefit ratio), sertraline (good risk-benefit ratio), citalopram/escitalopram (good risk-benefit ratio), orlistat (low to moderate risk-benefit ratio), and topiramate (moderate risk-benefit ratio). Atomoxetine has grade B evidence supporting its use.

## SAFETY SUMMARY

- Due to the potential for abuse, the stimulants are classified as Schedule II controlled substances. Atomoxetine, clonidine ER, guanfacine ER, and viloxazine ER are not classified as controlled substances.
- Various stimulants are contraindicated for use in patients with advanced arteriosclerosis, symptomatic CV disease, moderate to severe hypertension, hyperthyroidism, hypersensitivity to sympathomimetic amines, glaucoma, agitated states, history of drug abuse, tics, and in those using monoamine oxidase inhibitors (MAOIs). The stimulants carry a boxed warning for potential drug abuse and dependence. They also have warnings for increased risks of serious CV reactions, psychiatric AEs, suppression of growth, seizures, visual disturbance, peripheral vasculopathy, and priapism. Amphetamines have a warning for risk of serotonin syndrome when used in combination with other drugs affecting the serotonergic neurotransmitter systems.
  - Common AEs of stimulants include anorexia, decreased weight, tachycardia, anxiety, irritability, and insomnia.
  - Refer to the prescribing information for details on warnings, precautions, and AEs for individual products. For example:
    - QuilliChew ER can be harmful to patients with phenylketonuria (PKU) since it contains phenylalanine.
    - Because Concerta and Relexii tablets are nondeformable and do not appreciably change in shape in the gastrointestinal tract, they should not ordinarily be administered to patients with preexisting severe gastrointestinal narrowing.
    - The use of Daytrana may result in chemical leukoderma and contact sensitization; in addition, exposure of the application site to external heat sources should be avoided due to increased absorption of the drug.
    - Adhansia XR capsules contain FD&C yellow No. 5 dye (tartrazine), which may cause allergic-type reactions in susceptible patients.
- Atomoxetine is contraindicated for use in patients with narrow angle glaucoma, pheochromocytoma, severe CV disorders, hypersensitivity to any component of the product, and in those taking MAOIs. It carries a boxed warning for a rare increased risk of suicidal ideation in children and adolescents. It also has warnings for serious CV events, effects on blood pressure and heart rate, effects on growth, psychiatric AEs, rare cases of severe liver injury, urinary retention, and priapism.
  - Common AEs associated with atomoxetine include somnolence, nausea, and vomiting.
- Viloxazine ER is contraindicated with concurrent use of MAOIs and sensitive CYP1A2 substrates or CYP1A2 substrates with a narrow therapeutic index. Viloxazine ER carries a boxed warning for suicidal thoughts and behavior in children and adolescents. It also has warnings for effects on heart rate and blood pressure and the potential for somnolence and fatigue. Patients should be screened for bipolar disorder prior to use of viloxazine ER due to the risk of activation of mania or hypomania.
  - Common AEs associated with viloxazine ER include somnolence, nausea, and vomiting.
- The alpha<sub>2</sub>-adrenergic agonists are contraindicated in patients known to be hypersensitive to any constituent of the product. They carry warnings for increased risk of hypotension, bradycardia, and syncope; sedation and somnolence; rebound hypertension; and cardiac conduction abnormalities.
  - Common AEs associated with clonidine ER include somnolence, fatigue, and irritability while common AEs with guanfacine ER include somnolence, fatigue, and hypotension.

## DOSING AND ADMINISTRATION

**Table 4. Dosing and Administration**

Data as of November 22, 2021 HJI-U/CD-U/AVD

Page 11 of 23

This information is considered confidential and proprietary to OptumRx. It is intended for internal use only and should be disseminated only to authorized recipients. The contents of the therapeutic class overviews on this website ("Content") are for informational purposes only. The Content is not intended to be a substitute for professional medical advice, diagnosis, or treatment. Patients should always seek the advice of a physician or other qualified health provider with any questions regarding a medical condition. Clinicians should refer to the full prescribing information and published resources when making medical decisions.

Drug	Duration of action*	Available Formulations	Route	Usual Recommended Frequency	Comments
<b>Stimulants</b>					
Evekeo (amphetamine)	4 to 6 h	Tablets	Oral	<i>ADHD, narcolepsy</i> : Daily up to divided doses daily  <i>Exogenous obesity</i> : Divided doses daily	<i>ADHD and narcolepsy</i> The first dose should be given upon awakening; additional doses at intervals of 4 to 6 hours.
Evekeo ODT (amphetamine)	4 to 6 h	Orally disintegrating tablets	Oral	Once or twice daily in the morning	As soon as the blister pack is opened, the tablet should be placed on the patient's tongue and allowed to disintegrate without chewing or crushing. The tablet will disintegrate in saliva so that it can be swallowed.
Adzenys XR-ODT (amphetamine ER)	10 to 12 h	Orally disintegrating tablets	Oral	Daily in the morning	As soon as the blister pack is opened, the tablet should be placed on the patient's tongue and allowed to disintegrate without chewing or crushing. The tablet will disintegrate in saliva so that it can be swallowed.
Dyanavel XR (amphetamine ER)	Up to 13 h	Suspension, ER tablets	Oral	Daily in the morning	The bottle should be shaken before administration.  ER tablets may be chewed or swallowed whole. The 5 mg tablet may be split along the score line.
Adderall (mixed amphetamine salts)	4 to 6 h	Tablets	Oral	<i>ADHD, narcolepsy</i> : Daily up to divided doses daily	The first dose should be given on awakening, then additional doses at intervals of 4 to 6 hours.

Drug	Duration of action*	Available Formulations	Route	Usual Recommended Frequency	Comments
Adderall XR (mixed amphetamine salts ER)	10 to 12 h	Capsules	Oral	Daily in the morning	Capsules may be taken whole, or the capsule may be opened and the entire contents sprinkled on applesauce and consumed immediately. The dose of a single capsule should not be divided.
Mydayis (mixed amphetamine salts ER)	16 h	Capsules	Oral	Daily in the morning	Dosage adjustment is needed for severe renal impairment. Use in end stage renal disease (ESRD) is not recommended.  Capsules may be taken whole, or the capsule may be opened and the entire contents sprinkled on applesauce and consumed immediately in its entirety without chewing. The dose of a single capsule should not be divided.
Focalin (dexmethylphenidate)	3 to 5 h	Tablets	Oral	Twice daily	Separate doses by at least 4 hours.
Focalin XR (dexmethylphenidate ER)	8 to 12 h	Capsules	Oral	Daily in the morning	ER capsules may be taken whole, or the capsule may be opened and the entire contents sprinkled on applesauce and consumed immediately in its entirety without chewing. The dose of a single capsule should not be divided.



Drug	Duration of action*	Available Formulations	Route	Usual Recommended Frequency	Comments
ProCentra, Zenzedi (dextroamphetamine)	4 to 6 h	Solution (ProCentra) Tablets (Zenzedi)	Oral	<u>ADHD, narcolepsy</u> : Daily up to divided doses daily	The first dose should be given upon awakening; additional doses at intervals of 4 to 6 hours
Dexedrine Spansule (dextroamphetamine SR)	6 to 8 h	Capsules	Oral	<u>ADHD</u> Daily or twice daily  <u>Narcolepsy</u> Daily	
Vyvanse (lisdexamfetamine)	10 to 12 h	Capsules, chewable tablets	Oral	<u>ADHD, BED</u> : Daily in the morning	Dosage adjustment is needed for renal impairment/ESRD.  The capsules may be swallowed whole or can be opened, emptied, and mixed with yogurt, water, or orange juice and consumed immediately. A single capsule should not be divided.  The chewable tablets must be chewed thoroughly before swallowing. A single dose should not be divided.
Desoxyn (methamphetamine)	4 to 5 h	Tablets	Oral	Daily to twice daily	
Methylin, Ritalin (methylphenidate)	3 to 5 h	Chewable tablets, tablets (Ritalin), solution (Methylin)	Oral	Twice daily to 3 times daily	The chewable tablets should be taken with at least 8 ounces (a full glass) of water or other fluid.  The liquid and chewable tablets

Drug	Duration of action*	Available Formulations	Route	Usual Recommended Frequency	Comments
Methylphenidate ER	8 h	Tablets			<p>should be given 30 to 45 minutes before meals.</p> <p>The ER tablets may be used in place of the IR tablets when the 8-hour dosage of the ER product corresponds to the titrated 8-hour dosage of the IR products.</p> <p>The ER tablets must be swallowed whole and never crushed or chewed.</p>
Adhansia XR (methylphenidate ER)	13 h	Capsules	Oral	Daily in the morning	<p>The capsules may be taken whole or they can be opened and sprinkled onto applesauce or yogurt; the entire contents of the mixture should be consumed within 10 minutes, and should not be chewed.</p> <p>The dose of a single capsule should not be divided.</p>
Aptensio XR (methylphenidate ER)	12 h	Capsules	Oral	Daily in the morning	<p>The capsules may be taken whole or they can be opened and sprinkled onto applesauce; the applesauce should be consumed immediately and it should not be chewed.</p> <p>The dose of a single capsule should not be divided.</p>

Drug	Duration of action*	Available Formulations	Route	Usual Recommended Frequency	Comments
Concerta (methylphenidate ER)	12 h	Tablets	Oral	Daily in the morning	The tablets should not be chewed or crushed.
Methylphenidate ER					Note: An FDA analysis of methylphenidate ER products manufactured by UCB/Kremers (formerly Kudco) and Mallinckrodt indicated that in some individuals, they may deliver the drug in the body at a slower rate during the 7- to 12-hour range. As a result, the FDA changed the therapeutic equivalence of these products from AB to BX. Because these manufacturers have subsequently failed to demonstrate that their products are bioequivalent to the brand-name reference drug, the FDA proposed to withdraw their approval ( <i>FDA 2016</i> ).
Cotempla XR-ODT (methylphenidate ER)	12 h	Orally disintegrating tablets	Oral	Daily in the morning	As soon as the blister pack is opened, the tablet should be placed on the patient's tongue and allowed to disintegrate without chewing or crushing. The tablet will disintegrate in saliva so that it can be swallowed.
Jornay PM (methylphenidate ER)	10 h	Capsules	Oral	Daily in the evening	The capsule may be swallowed whole

Drug	Duration of action*	Available Formulations	Route	Usual Recommended Frequency	Comments
					or it may be opened and the contents sprinkled onto applesauce and given immediately. The capsule contents must not be crushed or chewed, the dose of a single capsule should not be divided, and the contents of the entire capsule should be taken at the same time.
Methylphenidate ER (CD)	6 to 9 h	Capsules	Oral	Daily in the morning	The capsule may be swallowed whole or it may be opened and the contents sprinkled onto a small amount (one tablespoon) of applesauce and given immediately, followed by some fluids. The capsule contents must not be crushed or chewed.
QuilliChew ER (methylphenidate ER)	8 h	Chewable tablets	Oral	Daily in the morning	A 10 mg or 15 mg dose can be achieved by breaking in half the functionally scored 20 mg and 30 mg tablets, respectively.
Quillivant XR (methylphenidate ER)	12 h	Suspension	Oral	Daily in the morning	The bottle of Quillivant XR should be shaken vigorously for 10 seconds prior to administration.  The suspension is stable for up to 4 months once reconstituted.

Drug	Duration of action*	Available Formulations	Route	Usual Recommended Frequency	Comments
Relexxii (methylphenidate ER 72 mg)	12 h	Tablet	Oral	Daily in the morning	The tablet must be swallowed whole with the aid of liquids, and must not be chewed, divided, or crushed.
Ritalin LA (methylphenidate ER)	6 to 9 h	Capsules	Oral	Daily in the morning	The capsule may be swallowed whole or may be administered by sprinkling the capsule contents on a small amount of applesauce; the contents should not be crushed, chewed, or divided. The mixture should be consumed immediately.
Daytrana (methylphenidate transdermal system)	12 h	Transdermal system	Transdermal	The patch should be applied 2 hours before an effect is needed and removed within 9 hours. It may be removed earlier than 9 hours if a shorter duration of effect is desired or late day side effects appear.	
Azstarys (serdexmethylphenidate/ dexmethylphenidate)	10 to 13 h	Capsules	Oral	Daily in the morning	The capsule may be swallowed whole or may be administered by sprinkling the capsule contents over 2 tablespoons of applesauce or 50 mL of water. The mixture should be consumed immediately.
<b>Non-stimulants</b>					
Strattera (atomoxetine)	At least 10 to 12 h	Capsules	Oral	Daily in the morning or divided dose in the morning and	Dosage adjustment is recommended for patients with moderate or severe



Drug	Duration of action*	Available Formulations	Route	Usual Recommended Frequency	Comments
				late afternoon/ early evening	hepatic insufficiency, for use with strong CYP2D6 inhibitors, and for patients known to be CYP2D6 poor metabolizers.  The capsules are not intended to be opened and should be taken whole.
Kapvay (clonidine ER)	At least 10 to 12 h	Tablets	Oral	Daily at bedtime or twice daily divided doses	With twice daily dosing, either an equal or higher split dosage should be given at bedtime.  The tablets should not be crushed, chewed, or broken prior to swallowing.  The initial dosage should be based on the degree of renal impairment.
Intuniv (guanfacine ER)	At least 8 to 12 h	Tablets	Oral	Daily in the morning or evening	The tablets should not be crushed, chewed, or broken prior to swallowing; they should not be administered with high fat meals, due to increased exposure.  It may be necessary to reduce the dosage in patients with significant renal and hepatic impairment.
Qelbree (viloxazine ER)	Throughout the day (specific duration not reported)	Capsule	Oral	Daily	The capsule may be swallowed whole or may be administered by sprinkling the capsule contents

Drug	Duration of action*	Available Formulations	Route	Usual Recommended Frequency	Comments
					over a teaspoon of applesauce. The mixture should be consumed within 2 hours, without chewing.

See the current prescribing information for full details

\*References: Prescribing information for individual products, *Medical Letter 2020*, *Pharmacist's Letter 2021*, *Krull 2020*.

## CONCLUSION

- Both CNS stimulants and nonstimulants may be used for the treatment of ADHD. In general, stimulants are first-line treatment due to their superior efficacy. Clinical evidence suggests that methylphenidate and amphetamines are equally efficacious, but some patients may respond to one stimulant and not the other. Various short-, intermediate- and long-acting formulations (eg, tablets/capsules, chewable/orally disintegrating tablets, solution/suspension, transdermal patch) are available to provide a range of dosing options. Although nonstimulants such as atomoxetine and alpha<sub>2</sub>-adrenergic agonists have smaller effect sizes, they may be used in patients who have failed or are intolerant to stimulants or when there is concern about possible abuse or diversion. The efficacy of the nonstimulant viloxazine ER in comparison to other nonstimulants is unknown. The alpha<sub>2</sub>-adrenergic agonists are approved both as monotherapy and as adjunctive therapy to stimulants, and they have been shown to improve both tic and ADHD symptoms in patients with comorbid tic disorder.
  - Current consensus clinical guidelines for the treatment of children and adolescents with ADHD recommend that stimulants are highly effective for reducing core symptoms of ADHD in children (*Wolraich et al 2019*).
- Ultimately, the choice of the initial agent for treatment of ADHD depends upon various factors such as: duration of desired coverage; ability of the child to swallow pills; coexisting tic disorder (use of alpha<sub>2</sub>-adrenergic agonists may be warranted); potential AEs, history of substance abuse in the patient or household member (eg, avoid stimulants or use stimulants with less potential for abuse [eg, lisdexamfetamine, osmotic-release preparation, methylphenidate patch]); and preference of the patient and parent/guardian (*Krull 2021a*).
- Various stimulants are indicated for treatment of narcolepsy and are generally considered to be second-line agents after modafinil/armodafinil due to their sympathomimetic AEs (*Scammell 2021*).
- Lisdexamfetamine is the only FDA-approved drug indicated for the treatment of moderate to severe BED, with demonstrated efficacy in reduction of mean binge days per week vs placebo. Direct comparison trials between lisdexamfetamine and other drugs used off-label to treat BED are lacking.

## REFERENCES

- Adderall [package insert], Horsham, PA: Teva Pharmaceuticals USA, Inc.; March 2020.
- Adderall XR [package insert], Lexington, MA: Takeda Pharmaceuticals America, Inc.; August 2020.
- Adhansia XR FDA clinical review. Food and Drug Administration website. February 7, 2019. [https://www.accessdata.fda.gov/drugsatfda\\_docs/nda/2019/212038Orig1s000MedR.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/nda/2019/212038Orig1s000MedR.pdf). Accessed November 19, 2021.
- Adhansia XR [package insert], Wilson, NC: Purdue Pharmaceuticals L.P.; June 2021.
- Adzenys XR-ODT [package insert], Grand Prairie, TX: Neos Therapeutics, Inc.; February 2018.
- Aigner M, Treasure J, Kaye W, Kasper S; WFSBP Task Force On Eating Disorders. World Federation of Societies of Biological Psychiatry (WFSBP) guidelines for the pharmacological treatment of eating disorders. *World J Biol Psychiatry*. 2011;12(6):400-443.
- Aptensio XR [package insert], Coventry, RI: Rhodes Pharmaceuticals, L.P.; June 2021.
- Azstarys [package insert], Grand Rapids, MI: Corium, Inc.; June 2021.
- Barbaresi WJ, Campbell L, Diekroger EA, et al. Society for Developmental and Behavioral Pediatrics clinical practice guideline for the assessment and treatment of children and adolescents with complex attention-deficit/hyperactivity disorder. *J Dev Behav Pediatr*. 2020;41 Suppl 2S:S35-S57. doi:10.1097/DBP.0000000000000770
- Brownley KA, Berkman ND, Peat CM, et al. Binge-eating disorder in adults: A systematic review and meta-analysis. *Ann Intern Med*. 2016;165(6):409-420.
- Bukstein O. Attention deficit hyperactivity disorder in adults: Epidemiology, pathogenesis, clinical features, course, assessment, and diagnosis. UpToDate Web site. 2021. <http://www.uptodate.com>. Updated March 26, 2021. Accessed November 19, 2021.
- Castells X, Blanco-Silvente L, Cunill R. Amphetamines for attention deficit hyperactivity disorder (ADHD) in adults. *Cochrane Database Syst Rev*. 2018;8:CD007813. doi: 10.1002/14651858.CD007813.pub3.

- Centers for Disease Control and Prevention (CDC). Attention Deficit Hyperactivity Disorder (ADHD). National Center for Health Statistics. <https://www.cdc.gov/nchs/fastats/adhd.htm>. Reviewed June 11, 2021. Accessed November 19, 2021.
- Chan E, Fogler JM, Hammerness PG. Treatment of attention-deficit/hyperactivity disorder in adolescents: a systematic review. *JAMA*. 2016;315(18):1997-2008.
- Childress AC, Brams M, Cutler AJ, et al. The efficacy and safety of Evekeo, racemic amphetamine sulfate, for treatment of attention-deficit/hyperactivity disorder symptoms: a multicenter, dose-optimized, double-blind, randomized, placebo-controlled crossover laboratory classroom study. *J Child Adolesc Psychopharmacol*. 2015;25(5):402-414. doi: 10.1089/cap.2014.0176
- Childress AC, Cutler AJ, Marraffino A, et al. A randomized, double-blind, placebo-controlled study of HLD200, a delayed-release and extended-release methylphenidate, in children with attention-deficit/hyperactivity disorder: An evaluation of safety and efficacy throughout the day and across settings. *J Child Adolesc Psychopharmacol*. 2020;30(1):2-14.
- Childress AC, Kollins SH, Cutler AJ, Marraffino A, Sikes CR. Efficacy, safety, and tolerability of an extended-release orally disintegrating methylphenidate tablet in children 6-12 years of age with attention-deficit/hyperactivity disorder in the laboratory classroom setting. *J Child Adolesc Psychopharmacol*. 2017;27(1):66-74.
- Clinical Pharmacology Website. <https://www.clinicalkey.com/pharmacology/login>. Accessed November 19, 2021.
- Concerta [package insert], Titusville, NJ: Janssen Pharmaceuticals, Inc.; June 2021.
- Cotempla XR-ODT [package insert], Grand Prairie, TX: Neos Therapeutics Brands, LCC; June 2021.
- Cortese S, Adamo N, Del Giovane C, et al. Comparative efficacy and tolerability of medications for attention-deficit hyperactivity disorder in children, adolescents, and adults: a systematic review and network meta-analysis. *Lancet Psychiatry*. 2018;5(9):727-738. doi: 10.1016/S2215-0366(18)30269-4.
- Cunill R, Castells X, Tobias A, Capellà D. Atomoxetine for attention deficit hyperactivity disorder in the adulthood: a meta-analysis and meta-regression. *Pharmacoevidemiol Drug Saf*. 2013;22(9):961-969.
- Daytrana [package insert], Miami, FL: Noven Therapeutics, LLC; June 2021.
- Desoxyyn [package insert], Lebanon, NJ: Recordati Rare Diseases Inc.; March 2019.
- Dexedrine Spansule [package insert], Bridgewater, NJ: Amneal Pharmaceuticals LLC; March 2019.
- Drugs@FDA [database on the Internet], Rockville (MD): Food and Drug Administration (US), Center for Drug Evaluation and Research; 2021. Available from: <http://www.accessdata.fda.gov/scripts/cder/daf/index.cfm>. Accessed November 19, 2021.
- Drugs for ADHD. *Med Lett Drugs Ther*. 2020;62 (1590):9-16.
- Dyanavel XR [package insert], Monmouth Junction: Tris Pharma, Inc.; November 2021.
- Elliott J, Johnston A, Husereau D, et al. Pharmacologic treatment of attention deficit hyperactivity disorder in adults: A systematic review and network meta-analysis. *PLoS One*. 2020;15(10):e0240584. doi: 10.1371/journal.pone.0240584
- Evekeo [package insert], Atlanta, GA: Arbor Pharmaceuticals, LLC; October 2019.
- Evekeo ODT [package insert], Atlanta, GA: Arbor Pharmaceuticals, LLC; April 2021.
- Faraone SV, Biederman J, Spencer TJ, Aleardi M. Comparing the efficacy of medications for ADHD using meta-analysis. *MedGenMed*. 2006;8(4):4.
- Faraone SV, Buitelaar J. Comparing the efficacy of stimulants for ADHD in children and adolescents using meta-analysis. *Eur Child Adolesc Psychiatry*. 2010a;19(4):353-364.
- Faraone SV, Glatt SJ. A comparison of the efficacy of medications for adult attention-deficit/hyperactivity disorder using meta-analysis of effect sizes. *J Clin Psychiatry*. 2010b;71(6):754-763.
- Feldman HM, Reiff MI. Clinical practice. Attention deficit-hyperactivity disorder in children and adolescents. *N Engl J Med*. 2014;370(9):838-846.
- Focalin [package insert], East Hanover, NJ: Novartis Pharmaceuticals Corporation.; June 2021.
- Focalin XR [package insert], East Hanover, NJ: Novartis Pharmaceuticals Corporation.; June 2021.
- Food and Drug Administration. Methylphenidate Hydrochloride Extended Release Tablets (generic Concerta) made by Mallinckrodt and Kudco. FDA Web site. <https://www.fda.gov/drugs/drug-safety-and-availability/methylphenidate-hydrochloride-extended-release-tablets-generic-concerta-made-mallinckrodt-and-kudco>. Updated November 4, 2016. Accessed November 24, 2021.
- Frick G, Yan B, Adler LA. Triple-bead mixed amphetamine salts (SHP465) in adults with ADHD: results of a phase 3, double-blind, randomized, forced-dose trial. *J Atten Disord*. 2020;24(3):402-413. doi: 10.1177/1087054717696771
- Garvey WT, Mechanick JI, Brett EM, et al; Reviewers of the AACE/ACE Obesity Clinical Practice Guidelines. American Association of Clinical Endocrinologists and American College of Endocrinology comprehensive clinical practice guidelines for medical care of patients with obesity. *Endocr Pract*. 2016;22 Suppl 3:1-203. doi: 10.4158/EP161365.GL.
- Gasior M, Hudson J, Quintero J, Ferreira-Cornwell MC, Radewonuk J, McElroy SL. A phase 3, multicenter, open-label, 12-month extension safety and tolerability trial of lisdexamfetamine dimesylate in adults with binge eating disorder. *J Clin Psychopharmacol*. 2017;37(3):315-322.
- Ghaderi A, Odeberg J, Gustafsson S, et al. Psychological, pharmacological, and combined treatments for binge eating disorder: a systematic review and meta-analysis. *PeerJ*. 2018;6:e5113. doi:10.7717/peerj.5113.
- Greenhill L, Kollins S, Abikoff H, et al. Efficacy and safety of immediate-release methylphenidate treatment for preschoolers with ADHD. *J Am Acad Child Adolesc Psychiatry*. 2006;45(11):1284-1293.
- Hirota T, Schwartz S, Correll CU. Alpha-2 agonists for attention-deficit/hyperactivity disorder in youth: a systematic review and meta-analysis of monotherapy and add-on trials to stimulant therapy. *J Am Acad Child Adolesc Psychiatry*. 2014;53(2):153-173.
- Intuniv [package insert], Lexington, MA: Shire US Inc.; August 2020.
- Hudson JI, McElroy SL, Ferreira-Cornwell MC, Radewonuk J, Gasior M. Efficacy of lisdexamfetamine in adults with moderate to severe binge-eating disorder: A randomized clinical trial. *JAMA Psychiatry*. 2017;74(9):903-910.
- Jadad AR, Boyle M, Cunningham C, Kim M, Schachar R. Treatment of attention-deficit/hyperactivity disorder. *Evid Rep Technol Assess (Summ)*. 1999;(11):i-viii, 1-341.
- Jornay PM [package insert], Cherry Hill, NJ: Ironshore Pharmaceuticals, Inc.; June 2021.
- Joseph A, Ayyagari R, Xie M, et al. Comparative efficacy and safety of attention-deficit/hyperactivity disorder pharmacotherapies, including guanfacine extended release: a mixed treatment comparison. *Eur Child Adolesc Psychiatry*. 2017;26(8):875-897.
- Kapvay [package insert], St. Michael, Barbados: Concordia Pharmaceuticals, Inc.; February 2020.

- Krull KR. Attention deficit hyperactivity disorder in children and adolescents: Clinical features and diagnosis. UpToDate Web site. 2019. <http://www.uptodate.com>. Updated November 27, 2019. Accessed November 19, 2021.
- Krull KR. Attention deficit hyperactivity disorder in children and adolescents: Epidemiology and pathogenesis. UpToDate Web site. 2021a. <http://www.uptodate.com>. Updated October 29, 2021. Accessed November 19, 2021.
- Krull KR. Attention deficit hyperactivity disorder in children and adolescents: Overview of treatment and prognosis. UpToDate Web site. 2020. <http://www.uptodate.com>. Updated February 3, 2020. Accessed November 19, 2021.
- Krull KR. Attention deficit hyperactivity disorder in children and adolescents: Treatment with medications. UpToDate Web site. 2021b. <http://www.uptodate.com>. Updated May 13, 2021. Accessed November 19, 2021.
- Krull KR. Pharmacology of drugs used to treat attention deficit hyperactivity disorder in children and adolescents. UpToDate Web site. 2021c. <http://www.uptodate.com>. Updated May 13, 2021. Accessed November 19, 2021.
- Maski K, Trotti LM, Kotagal S, et al. Treatment of central disorders of hypersomnolence: an American Academy of Sleep Medicine clinical practice guideline. *J Clin Sleep Med*. 2021;17(9):1881-1893. doi:10.5664/jcsm.9328
- McElroy SL, Hudson J, Ferreira-Cornwell MC, Radewonuk J, Whitaker T, Gasior M. Lisdexamfetamine dimesylate for adults with moderate to severe binge eating disorder: Results of two pivotal phase 3 randomized controlled trials. *Neuropsychopharmacology*. 2016;41(5):1251-1260.
- Methylin oral solution [package insert], Florham Park, NJ: Shionogi Inc.; July 2021.
- Methylphenidate chewable tablets [package insert], Central Islip, NY: Ascent Pharmaceuticals, Inc.; August 2021.
- Methylphenidate ER [package insert], Newtown, PA: KVK-Tech, Inc.; April 2021.
- Methylphenidate ER (CD) [package insert], Philadelphia, PA: Lannett Company, Inc.; May 2021.
- Methylphenidate hydrochloride extended-release tablets (generic Concerta) made by Mallinckrodt and Kudco. FDA Web site. November 4, 2016.
- Murphy TK, Lewin AB, Storch EA, Stock S; American Academy of Child and Adolescent Psychiatry (AACAP) Committee on Quality Issues (CQI). Practice parameter for the assessment and treatment of children and adolescents with tic disorders. *J Am Acad Child Adolesc Psychiatry*. 2013;52(12):1341-1359.
- Mydayis [package insert], Lexington, MA: Shire US Inc.; August 2020.
- Nasser A, Liranso T, Adewole T, et al. A Phase III, randomized, placebo-controlled trial to assess the efficacy and safety of once-daily SPN-812 (viloxazine extended-release) in the treatment of attention-deficit/hyperactivity disorder in school-age children. *Clin Ther*. 2020;42(8):1452-1466.
- Nasser A, Liranso T, Adewole T, et al. A phase 3, placebo-controlled trial of once-daily viloxazine extended-release capsules in adolescents with attention-deficit/hyperactivity disorder. *J Clin Psychopharmacol*. 2021a;41(4):370-380. doi:10.1097/JCP.0000000000001404
- Nasser A, Liranso T, Adewole T, et al. Once-daily SPN-812 200 and 400 mg in the treatment of ADHD in school-aged children: A Phase III randomized, controlled trial. *Clin Ther*. 2021b;S0149-2918(21)00054-0.
- Newcorn JH, Kratochvil CJ, Allen AJ, et al; Atomoxetine/Methylphenidate Comparative Study Group. Atomoxetine and osmotically released methylphenidate for the treatment of attention deficit hyperactivity disorder: acute comparison and differential response. *Am J Psychiatry*. 2008;165(6):721-730.
- Orange Book: Approved Drug Products with Therapeutic Equivalence Evaluations. Food and Drug Administration Web site. <http://www.accessdata.fda.gov/scripts/cder/ob/default.cfm>. Accessed November 19, 2021.
- Osland ST, Steeves TD, Pringsheim T. Pharmacological treatment for attention deficit hyperactivity disorder (ADHD) in children with comorbid tic disorders. *Cochrane Database Syst Rev*. 2018;6:CD007990. doi:10.1002/14651858.CD007990.pub3.
- Padilha SCOS, Virtuoso S, Tonin FS, Borba HHL, Pontarolo R. Efficacy and safety of drugs for attention deficit hyperactivity disorder in children and adolescents: a network meta-analysis. *Eur Child Adolesc Psychiatry*. 2018;27(10):1335-1345. doi: 10.1007/s00787-018-1125-0.
- PL Detail-Documents, Comparison of ADHD medications. *Pharmacist's Letter/Prescriber's Letter*. September 2021.
- Pliszka SR, Wilens TE, Bostrom S, et al. Efficacy and safety of HLD200, delayed-release and extended-release methylphenidate, in children with attention-deficit/hyperactivity disorder. *J Child Adolesc Psychopharmacol*. 2017;27(6):474-482. doi: 10.1089/cap.2017.0084.
- Practice parameter for the assessment and treatment of children and adolescents with tic disorders. *J Am Acad Child Adolesc Psychiatry*. 2013;52(12):1341-1359.
- Pringsheim T, Okun MS, Müller-Vahl K, et al. Practice guideline recommendations summary: Treatment of tics in people with Tourette syndrome and chronic tic disorders. *Neurology*. 2019;92(19):896-906.
- ProCentra [package insert], Newport, KY: Independence Pharmaceuticals, LLC; February 2017.
- Punja S, Shamsseer L, Hartling L, et al. Amphetamines for attention deficit hyperactivity disorder (ADHD) in children and adolescents. *Cochrane Database Syst Rev*. 2016;2:CD009996.
- Qelbree [package insert], Rockville, MD: Supernus Pharmaceuticals, Inc.; April 2021.
- QuilliChew ER [package insert], Monmouth Junction, NJ: Tris Pharma, Inc.; June 2021.
- Quillivant XR [package insert], Monmouth Junction, NJ: Tris Pharma, Inc.; June 2021.
- Relexxii [package insert], Bridgewater, NJ: Vertical Pharmaceuticals, LLC; July 2021.
- Rodrigues R, Lai MC, Beswick A, et al. Practitioner Review: Pharmacological treatment of attention-deficit/hyperactivity disorder symptoms in children and youth with autism spectrum disorder: a systematic review and meta-analysis. *J Child Psychol Psychiatry*. 2021;62(6):680-700. doi:10.1111/jcpp.13305
- Ritalin [package insert], East Hanover, NJ: Novartis Pharmaceuticals Corporation; June 2021.
- Ritalin LA [package insert], East Hanover, NJ: Novartis Pharmaceuticals Corporation; June 2021.
- Scammell TE. Treatment of narcolepsy in adults. UpToDate Web site. 2021. <http://www.uptodate.com>. Updated April 30, 2021. Accessed November 11, 2021.
- Schneider E, Higgs S, Dourish CT. Lisdexamfetamine and binge-eating disorder: A systematic review and meta-analysis of the preclinical and clinical data with a focus on mechanism of drug action in treating the disorder [published online ahead of print. *Eur Neuropsychopharmacol*. 2021;53:49-78. doi:10.1016/j.euroneuro.2021.08.001
- Schwartz S, Correll CU. Efficacy and safety of atomoxetine in children and adolescents with attention-deficit/hyperactivity disorder: results from a comprehensive meta-analysis and metaregression. *J Am Acad Child Adolesc Psychiatry*. 2014;53(2):174-187.

- Storebø OJ, Ramstad E, Krogh HB, et al. Methylphenidate for children and adolescents with attention deficit hyperactivity disorder (ADHD). *Cochrane Database Syst Rev*. 2015;11:CD009885.
- Strattera [package insert], Indianapolis, IN: Lilly USA, Inc.; February 2020.
- Stuhec M, Lukić P, Locatelli I. Efficacy, acceptability, and tolerability of lisdexamfetamine, mixed amphetamine salts, methylphenidate, and modafinil in the treatment of attention-deficit hyperactivity disorder in adults: a systematic review and meta-analysis. *Ann Pharmacother*. 2019;53(2):121-133. doi:10.1177/1060028018795703.
- Stuhec M, Munda B, Svab V, Locatelli I. Comparative efficacy and acceptability of atomoxetine, lisdexamfetamine, bupropion and methylphenidate in treatment of attention deficit hyperactivity disorder in children and adolescents: a meta-analysis with focus on bupropion. *J Affect Disord*. 2015;178:149-159.
- Vyvanse [package insert], Lexington, MA: Takeda Pharmaceuticals America, Inc.; July 2021.
- Weisler RH, Greenbaum M, Arnold V, et al. Efficacy and safety of SHP465 mixed amphetamine salts in the treatment of attention-deficit/hyperactivity disorder in adults: results of a randomized, double-blind, placebo-controlled, forced-dose clinical study. *CNS Drugs*. 2017;31(8):685-697.
- Wigal T, Brams M, Frick G, Yan B, Madhoo M. A randomized, double-blind study of SHP465 mixed amphetamine salts extended-release in adults with ADHD using a simulated adult workplace design. *Postgrad Med*. 2018a;130(5):481-493. doi: 10.1080/00325481.2018.1481712
- Wigal T, Childress A, Frick G, Yan B, Wigal S, Madhoo M. Effects of SHP465 mixed amphetamine salts in adults with ADHD in a simulated adult workplace environment. *Postgrad Med*. 2018b;130(1):111-121. doi: 10.1080/00325481.2018.1389227
- Wigal T, Lopez F, Frick G, Yan B, Robertson B, Madhoo M. A randomized, double-blind, 3-way crossover, analog classroom study of SHP465 mixed amphetamine salts extended-release in adolescents with ADHD. *Postgrad Med*. 2019;131(3):212-224. doi: 10.1080/00325481.2019.1574402
- Wigal SB, Wigal T, Childress A, Donnelly GAE, Reiz JL. The time course of effect of multilayer-release methylphenidate hydrochloride capsules: a randomized, double-blind study of adults with ADHD in a simulated adult workplace environment. *J Atten Disord*. 2020;24(3):373-383. doi: 10.1177/1087054716672335
- Wolraich ML, Hagan JF Jr., Allan C, et al; Subcommittee on Children and Adolescents with Attention-Deficit/Hyperactive Disorder. Clinical practice guideline for the diagnosis, evaluation, and treatment of attention-deficit/hyperactivity disorder in children and adolescents. *Pediatrics*. 2019;144(4):e20192528. doi: 10.1542/peds.2019-2528
- Yager J, Devlin MF, Halmi KA, et al. Guideline watch (August 2012): Practice guideline for the treatment of patients with eating disorders, 3rd edition. Psychiatry Online Web site. <http://psychiatryonline.org/guidelines>. Accessed November 19, 2021.
- Yager J, Devlin MF, Halmi KA, et al. Practice guideline for the treatment of patients with eating disorders, 3rd edition (2006). Psychiatry Online Web site. <http://psychiatryonline.org/guidelines>. Accessed November 19, 2021.
- Zenedi [package insert], Atlanta, GA: Arbor Pharmaceuticals, LLC; March 2021.

Publication Date: December 9, 2021