

Therapeutic Class Overview

Antivirals, Influenza

INTRODUCTION

- Influenza is an infectious respiratory illness caused by the influenza A and influenza B viruses. Influenza epidemics occur annually in the United States, typically from late fall to early spring. Although the majority of infected individuals recover without complications, some cases of influenza result in severe illness or death (*Grohskopf et al 2017*).
- The virus is primarily transmitted through direct contact large-particle respiratory droplets from an infected individual's coughs and sneezes. It is also spread through contact with surfaces contaminated by infected respiratory droplets. Adults begin to shed virus 1 day prior to symptom onset, and they remain contagious for 5 to 7 days after falling ill (*Centers for Disease Control and Prevention [CDC] 2016[a]*).
- Signs and symptoms of uncomplicated influenza illness include fever, myalgia, headache, malaise, nonproductive cough, sore throat, and rhinitis. Complications of influenza infection include sinusitis, otitis media, pneumonia, sepsis, and exacerbation of chronic medical conditions. Elderly adults, young children, pregnant women, and patients with chronic medical conditions have a higher risk of developing complications from influenza (*CDC 2016[b]*).
- Annual influenza vaccination is the most effective method for preventing seasonal influenza virus infection and its complications. Antiviral prescription medications are also available for influenza prophylaxis and treatment; however, antiviral chemoprophylaxis is not a substitute for annual influenza vaccination (*Grohskopf et al 2017*).
- Initiation of antiviral therapy to treat influenza is recommended as early as possible for patients with confirmed or suspected influenza who are hospitalized, have severe, complicated, or progressive illness, or are at higher risk for influenza complications (*Fiore et al 2011*). Additionally, due to the increased influenza activity and a lower vaccine effectiveness for the 2017-2018 influenza season, a December 2017 CDC advisory recommends that all hospitalized patients and all high-risk patients (hospitalized or outpatient) with suspected influenza should be treated as soon as possible with a neuraminidase inhibitor. Although initiation within 2 days of symptom onset is ideal, the CDC is stating that benefit may still be seen even when treatment is initiated later (*CDC 2017*).
- Two classes of antiviral medications are available and will be reviewed. The adamantanes include amantadine and Flumadine (rimantadine). The neuraminidase inhibitors include Rapivab (peramivir), Relenza (zanamivir), and Tamiflu (oseltamivir).
- Although the adamantanes are active against influenza A virus, resistance is high amongst currently circulating virus strains. The adamantanes lack activity against influenza B virus. Therefore, amantadine and rimantadine are not recommended for treatment or chemoprophylaxis during the current influenza season (*CDC 2018*).
- The neuraminidase inhibitors are active against both influenza A and influenza B viruses. Rapivab (peramivir), Relenza (zanamivir), and oseltamivir are the only antivirals recommended for the current influenza season in the United States (*CDC 2018*).
- Circulating influenza viruses are constantly evolving, and drug-resistant influenza virus strains have been reported. Prescribers should refer to influenza drug susceptibility patterns and treatment effects when selecting an antiviral agent (*CDC 2018*).
- Medispan class: Antiparkinson, Dopaminergic and Influenza Agents. The only agent from the Antiparkinson, Dopaminergic category that will be included in this review is amantadine for the influenza indication.

Table 1. Medications Included Within Class Review

Drug	Generic Availability
amantadine	✓
Flumadine (rimantadine)	✓
Rapivab (peramivir)	-
Relenza (zanamivir)	-
Tamiflu (oseltamivir)	✓

(*Drugs @FDA 2018, Orange Book: Approved Drug Products with Therapeutic Equivalence Evaluations 2018*)

INDICATIONS

Table 2. Food and Drug Administration Approved Indications

Indication ¹	amantadine ²	Flumadine (rimantadine)	Rapivab ³ (peramivir)	Relenza ⁴ (zanamivir)	Tamiflu ⁵ (oseltamivir)
Prophylaxis and treatment of signs and symptoms of infection caused by various strains of influenza A virus	✓				
Prophylaxis and treatment of illness caused by various strains of influenza A virus in adults (17 years and older)		✓			
Prophylaxis against influenza A virus in children (1 to 16 years of age)		✓			
Treatment of acute uncomplicated influenza in patients 2 years and older who have been symptomatic for no more than 2 days			✓		
Prophylaxis of influenza in adults and pediatric patients aged 5 years and older				✓	
Treatment of uncomplicated acute illness due to influenza A and B virus in adults and pediatric patients aged 7 years and older who have been symptomatic for no more than 2 days				✓	
Prophylaxis of influenza A and B in patients 1 year and older					✓
Treatment of acute, uncomplicated illness due to influenza A and B infection in patients 2 weeks of age and older who have been symptomatic for no more than 48 hours					✓

¹ The changing of viruses over time is a limitation of use for antivirals. The emergence of resistance mutations could decrease drug effectiveness. Other factors, such as changes in viral virulence, may also diminish clinical benefit of antivirals. Prescribers should consider available information on influenza drug susceptibility patterns and treatment effects when selecting an antiviral.

² Amantadine is also indicated in the treatment of parkinsonism and drug-induced extrapyramidal reactions.

³ Limitations of use for Rapivab (peramivir):

- Efficacy is based on clinical trials of naturally occurring influenza in which the predominant influenza infections were influenza A virus; a limited number of subjects infected with influenza B virus were enrolled.
- Efficacy could not be established in patients with serious influenza requiring hospitalization.

⁴ Limitations of use for Relenza (zanamivir):

- Not recommended for treatment or prophylaxis of influenza in individuals with underlying airways disease (such as asthma or chronic obstructive pulmonary disease) due to the risk of serious bronchospasm.
- Has not been proven effective for treatment of influenza in individuals with underlying airways disease.
- Has not been proven effective for prophylaxis of influenza in the nursing home setting.

⁵ Limitations of use for Tamiflu (oseltamivir):

- Not recommended for patients with end-stage renal disease not undergoing dialysis.

(Prescribing information: amantadine capsules 2017, amantadine oral solution 2015, amantadine tablets 2017, Flumadine 2010, Rapivab 2018, Relenza 2016, Tamiflu 2018)

- Information on indications, mechanism of action, pharmacokinetics, dosing, and safety has been obtained from the prescribing information for the individual products, except where noted otherwise.

CLINICAL EFFICACY SUMMARY

Adamantanes

- Clinical trials have demonstrated that the adamantanes are effective in both the prophylaxis and treatment of influenza A virus (*Bryson et al 1980, Crawford et al 1988, Dolin et al 1982, Hall et al 1987, Hayden et al 1989, Jackson et al 2011, Jefferson et al 2006[a], Jefferson et al 2006[b], Monto et al 1995, Reuman et al 1989*).
- One systematic review assessed the efficacy and safety of adamantanes in healthy adults by analyzing 20 prophylaxis and 13 treatment randomized trials comparing amantadine or rimantadine with placebo. For prophylaxis, amantadine was 61% better than placebo at reducing influenza risk ($P < 0.001$). Although rimantadine was 72% better than placebo at preventing influenza, statistical significance was not achieved. There was significant heterogeneity between the prophylaxis trials, and only a small sample size was available for rimantadine compared to amantadine. For treatment, amantadine and rimantadine both reduced the duration of fever by one day. Both agents caused gastrointestinal side effects, but amantadine caused significantly more adverse effects in the central nervous system than rimantadine (*Jefferson et al 2006[a]*).
- Influenza A virus resistance to amantadine and rimantadine has developed over the years. During the 2009 to 2010 influenza season, 100% of the 18 influenza H3N2 viruses tested in the United States were resistant to adamantanes. Similarly, 99.8% of the pandemic H1N1 viruses tested were resistant to adamantanes. Due to influenza A virus resistance and lack of activity against influenza B virus, the adamantanes are not recommended for the current influenza season (*CDC 2010[b], CDC 2018*).

Neuraminidase inhibitors

- The neuraminidase inhibitors have demonstrated efficacy for their respective indications. Relenza (zanamivir) inhalation and oral oseltamivir are effective in both the prophylaxis and treatment of influenza A and B. Clinical trials have demonstrated a reduction in laboratory-confirmed influenza, illness, fever duration, secondary complications, and household contacts with influenza infection (*Aoki et al 2003, Chik et al 2004, Cooper et al 2003, Fry et al 2014, Halloran et al 2007, Hayden et al 1997, Hayden et al 1999, Hayden et al 2000, Hayden et al 2004, Hedrick et al 2000, Hiba et al 2011, Kaiser et al 2003, Kawai et al 2005, Kawai et al 2006, Lin et al 2006, MIST Study Group 1998, Monto et al 1999[a], Monto et al 1999[b], Monto et al 2002, Nicholson et al 2000, Peters et al 2001, Reuman et al 1989, Singh et al 2003, Treanor et al 2000, Turner et al 2003, Wang et al 2012, Welliver et al 2001, Whitley et al 2001*).
- One systematic review analyzed 20 oseltamivir and 26 Relenza (zanamivir) randomized, placebo-controlled trials in order to better define their efficacy and safety. In prophylaxis trials, the risk of symptomatic influenza was reduced by 3.05% in patients treated with oseltamivir compared to placebo and 1.98% in patients treated with Relenza (zanamivir) compared to placebo. In adults, the time to first alleviation of symptoms was reduced by 0.7 days ($P < 0.0001$) in patients receiving oseltamivir compared to placebo and 0.6 days ($P < 0.00001$) in patients receiving Relenza (zanamivir) compared to placebo. Oseltamivir significantly reduced the time to alleviation of symptoms in non-asthmatic children and decreased the incidence of self-reported pneumonia. Relenza (zanamivir) significantly reduced the risk of bronchitis in adults with influenza. Neither treatment was a significant improvement over placebo in time to symptom alleviation in asthmatic children or risk of hospitalizations, otitis media, or sinusitis. Many studies included were at a high risk of selection bias due to inadequate reporting and a high risk of attrition bias due to selective reporting. All trials were sponsored by the manufacturers (*Jefferson et al 2014*).
- In a systematic review of other published systematic reviews and meta-analyses, treatment of influenza with neuraminidase inhibitors (oseltamivir or zanamivir) was found to be likely effective in reducing mortality amongst hospitalized patients; the odds of mortality appeared especially lower when therapy was started early (≤ 48 hours of symptom onset). When used for treatment in the general population, these agents appear to reduce the duration of symptoms by approximately 0.5 to 1 day. Both oseltamivir and zanamivir were found likely to be effective at reducing secondary symptomatic influenza transmission when used prophylactically (*Doll et al 2017*).
- Rapivab (peramivir) intravenous (IV) infusion is approved for the treatment of influenza A and B in adults. The primary endpoint for the main clinical trial supporting Food and Drug Administration (FDA)-approval of Rapivab (peramivir) was time to alleviation of symptoms. The trial evaluated 296 previously healthy adults presenting with the onset of influenza-like illness within the previous 48 hours and a positive influenza rapid antigen test. In this multicenter, double-blind,

placebo-controlled clinical trial, patients were randomized to Rapivab (peramivir) 300 mg, 600 mg, or placebo as a single IV dose. Acetaminophen use was permitted. Patients self-reported body temperature, symptoms, and resumption of activities over 14 days. The primary endpoint, the median time to alleviation of symptoms, was significantly earlier with Rapivab (peramivir) 300 mg (59.1 hours) and 600 mg (59.9 hours) compared to placebo (81.8 hours; both $P=0.0092$). There was no significant difference in the incidence of all adverse events in patients receiving Rapivab (peramivir) compared to placebo. Diarrhea was the most common adverse event, occurring in 14.1%, 15.2% and 17% of the Rapivab (peramivir) 300 mg, 600 mg, and placebo groups, respectively (Kohno et al 2010).

- Although studies have evaluated Rapivab (peramivir) in hospitalized patients and in children, both of these populations are not included in the FDA-approved labeling (De Jong et al 2014, Ison et al 2014, Ison et al 2013, Sugaya et al 2012). The Phase 3 clinical trial of Rapivab (peramivir) in hospitalized influenza patients failed to meet its primary endpoint of reducing the time to clinical resolution compared to placebo. There are no clinical endpoints that have been validated for clinical trials of neuraminidase inhibitors treating hospitalized patients with influenza (FDA 2014). In 2009, the United States issued an Emergency Use Authorization (EUA) program allowing Rapivab (peramivir) for the treatment of suspected or confirmed 2009 H1N1 influenza A virus infection in hospitalized patients (Birnkranz 2009). Patients eligible for treatment were hospitalized, unable to tolerate or unresponsive to other available antivirals, or lacked a dependable oral or inhalation drug delivery route. The Public Health Emergency determination for the 2009 H1N1 influenza pandemic expired on June 23, 2010 (CDC 2010[a]).
- Numerous placebo-controlled trials have demonstrated the efficacy of neuraminidase inhibitors individually, but head-to-head trials directly comparing the agents are limited. One randomized, double-blind, placebo-controlled safety trial compared the use of oseltamivir, Relenza (zanamivir), and placebo in 390 healthy adults for influenza chemoprophylaxis over 16 weeks. The study showed that both treatments were well tolerated compared to placebo, and there were no discontinuations due to adverse events (Anekthananon et al 2013).
- A Phase 3 multinational, multicenter, double-blind, randomized, noninferiority trial compared a single dose of 300 or 600 mg IV Rapivab (peramivir) to 5 days of oral oseltamivir in 1,091 patients with seasonal influenza. The primary endpoint, time to alleviation of influenza symptoms, had a median of 78.0 hours in patients receiving 300 mg of Rapivab (peramivir), 81.0 hours in patients receiving 600 mg of Rapivab (peramivir), and 81.8 hours in patients receiving oseltamivir. Both strengths of Rapivab (peramivir) were noninferior to oseltamivir with a noninferiority margin of 0.170. There was no significant difference between treatments in the incidence of complications of influenza infection (Kohno et al 2011).
- A meta-analysis including 2 controlled clinical trials and 5 observational trials (N = 1676) examined the comparative efficacy of IV Rapivab (peramivir) and oral oseltamivir in the treatment of seasonal influenza. No significant differences between treatments were noted for the following outcomes: mortality, hospital length of stay, virus titer 48 hours after admission, and incidence of adverse events. However, the time to resolution of influenza symptoms or fever was shorter with Rapivab (peramivir) versus oseltamivir treatment (mean difference, -7.17 hours; $p < 0.01$) (Lee et al 2017).
- Observational studies comparing the clinical efficacy of Rapivab (peramivir), Relenza (zanamivir), and oseltamivir in treating influenza have demonstrated within-class variation in the time to alleviation of influenza symptoms. The lack of robust data from randomized, head-to-head trials prevents the recommendation of one neuraminidase inhibitor over another. Local and seasonal susceptibility trends, route of administration, and patient-specific factors such as age and compliance should be taken into account when selecting an agent for antiviral drug therapy (Kawai et al 2008, Takemoto et al 2013).
- While influenza virus strains resistant to specific neuraminidase inhibitors have emerged, overall resistance remains low. According to surveillance data on seasonal influenza virus strains, the rate of resistance to oseltamivir is 1 to 3% and resistance to Relenza (zanamivir) is less than 1% (Li et al 2015).

CLINICAL GUIDELINES

- Annual influenza vaccination is the most effective method for preventing seasonal influenza virus infection and its complications. All individuals six months of age and older should receive an influenza vaccination each year, unless contraindicated. The live attenuated intranasal influenza vaccine is not recommended during the 2017 to 2018 influenza season due to low effectiveness. The prophylactic antiviral administration is not a substitute for early influenza vaccination (Grohskopf et al 2017).
- Amantadine and rimantadine are not recommended for antiviral treatment or prophylaxis of influenza A virus strains in the United States due to high rates of resistance (American Academy of Pediatrics [AAP] 2017, Fiore et al 2011, CDC 2018).

- The antivirals recommended by the CDC for the current influenza season include oseltamivir, Relenza (zanamivir) and Rapivab (peramivir). Routine or widespread use of antivirals for chemoprophylaxis is not recommended due to concerns for viral resistance. Oseltamivir and Relenza (zanamivir) are recommended for post-exposure prophylaxis in patients who are severely immunosuppressed and in patients at a high risk for influenza complications who are either not a candidate for vaccination or received their annual vaccination less than 2 weeks prior to exposure (*CDC 2018*).
- Treatment of influenza with antiviral therapy is recommended as early as possible for patients with confirmed or suspected influenza who are hospitalized, have severe, complicated, or progressive illness, or are at a high risk for complications (*CDC 2018*).
- Populations at a high risk for influenza complications and recommended to receive antiviral treatment include children younger than 2 years old, adults age 65 and above, pregnant or postpartum women, American Indians, Alaska Natives, obese patients with a body mass index (BMI) of 40 kg/m² and above, patients younger than 19 years old receiving long-term treatment with aspirin, residents of nursing homes, and patients with immunosuppression, chronic disorders (eg, pulmonary, cardiovascular, renal, hepatic, hematological and metabolic), or neurologic conditions (*CDC 2018*). Additionally, due to the increased influenza activity and a lower vaccine effectiveness for the 2017-2018 influenza season, a December 2017 CDC advisory recommends that all hospitalized patients and all high-risk patients (hospitalized or outpatient) with suspected influenza should be treated as soon as possible with a neuraminidase inhibitor. Although initiation within 2 days of symptom onset is ideal, the CDC is stating that benefit may still be seen even when treatment is initiated later (*CDC 2017*).
- Antiviral therapy works best when administered within 48 hours of symptom onset. Treatment initiation should not be delayed for the results of diagnostic testing. Early administration of antivirals may shorten the duration of fever, reduce the risk of influenza-related complications such as otitis media and pneumonia, reduce death in hospitalized patients, and decrease the duration of hospitalization in hospitalized children (*CDC 2018*).

SAFETY SUMMARY

- Common adverse events with adamantanes include nausea, dizziness, insomnia, headache, anorexia, dry mouth, and agitation.
- Amantadine and rimantadine should be used with caution in patients with epilepsy due to an increased risk for seizures.
- Amantadine has anticholinergic effects and is contraindicated in patients with untreated angle closure glaucoma. There have also been reports of death from overdose and suicide attempts with amantadine.
- Common adverse events with neuraminidase inhibitors include nausea, vomiting, and headache. The most common adverse effect with Rapivab (peramivir) is diarrhea.
- All three neuraminidase inhibitors have labeled warnings for neuropsychiatric events such as hallucinations and delirium. Patients should be monitored for signs of abnormal behavior.
- Oseltamivir and Rapivab (peramivir) have warnings for serious skin and hypersensitivity reactions, including Stevens-Johnson Syndrome.
- Relenza (zanamivir) has a warning for bronchospasm and should not be used in patients with asthma or chronic obstructive pulmonary disease. It is also contraindicated in patients with milk protein allergies.

DOSING AND ADMINISTRATION

Table 3. Dosing and Administration*

Drug	Available Formulations	Route	Usual Recommended Frequency	Comments
amantadine	Capsules, oral solution, tablets	Oral	Once daily or twice daily <u>Adults:</u> 200 mg once daily or 100 mg twice daily <u>Pediatric patients:</u> 1 to 9 years: 4.4 to 8.8 mg/kg/day not to exceed 150 mg per day	Should be taken for 10 days following a known exposure. If using in conjunction with vaccine until antibody response, then take for 2 to 4 weeks. Treatment of illness should be started within 24 to 48 hours of symptom onset and continued

Drug	Available Formulations	Route	Usual Recommended Frequency	Comments
			<p>9 to 12 years: 100 mg twice daily</p> <p>The safety and efficacy of amantadine in newborn infants and infants below the age of 1 year have not been established.</p>	<p>for 24 to 48 hours after symptoms disappear.</p> <p>For adult patients intolerant to 200 mg daily dose because of central nervous system or other toxicities: 100 mg daily dose</p> <p>Because amantadine is primarily excreted in the urine, it accumulates in the plasma and in the body when renal function declines. Thus, the dose of amantadine should be reduced in patients with renal impairment and in individuals who are 65 years of age or older according to the following:</p> <p><u>For CrCl = 30 to 50 mL/min:</u> 200 mg 1st day, then 100 mg daily</p> <p><u>For CrCl = 15 to 29 mL/min:</u> 200 mg 1st day, then 100 mg on alternate days</p> <p><u>For CrCl < 15 mL/min and HD:</u> 200 mg every 7 days</p> <p><u>For patients ≥ 65 years:</u> 100 mg once daily</p> <p>The dose of amantadine may need reduction in patients with congestive heart failure, peripheral edema, or orthostatic hypotension.</p>
Flumadine (rimantadine)	Tablets	Oral	<p>Twice daily</p> <p>Adults (17 years and older) <u>Treatment:</u> 100 mg twice daily for 7 days</p> <p><u>Prophylaxis:</u> 100 mg twice daily</p> <p>Pediatric patients <u>Prophylaxis in patients 1 to 9 years:</u> 5 mg/kg/day, not to exceed 150 mg per day</p>	<p>Treatment of illness should be started within 48 hours of symptoms. A suspension can be made from the tablets and is stable for 14 days.</p> <p>Dose adjustment in patients ≥ 65 years: 100 mg once daily</p> <p>Dose adjustment in patients with CrCl < 29 mL/min: 100 mg daily</p> <p>Dose adjustment in patients with</p>

Drug	Available Formulations	Route	Usual Recommended Frequency	Comments
			<p><u>10 to 16 years</u>: Refer to the adult dose</p> <p>The safety and efficacy of rimantadine in pediatric patients below the age of 1 year have not been established.</p>	<p>severe hepatic dysfunction: 100 mg daily</p>
<p>Rapivab (peramivir)</p>	<p>Injection</p>	<p>IV</p>	<p><u>Patients ≥ 13 years</u>: 600 mg as a single dose</p> <p><u>Patients < 13 years</u>: 2 to 12 years: 12 mg/kg (maximum dose 600 mg) as a single dose</p> <p>Safety and effectiveness in pediatric patients < 2 years of age have not been established.</p>	<p>One time dose should be provided within 2 days of onset of influenza symptoms</p> <p>A single dose administered by IV infusion for a minimum of 15 minutes.</p> <p>Rapivab must be diluted prior to administration.</p> <p>Dose adjustment in adults and adolescents 13 years of age or older with CrCl = 30 to 49 mL/min: 200 mg</p> <p>Dose adjustment in pediatric patients 2 to 12 years of age with CrCl = 30 to 49 mL/min: 4 mg/kg</p> <p>Dose adjustment in adults and adolescents 13 years of age or older with CrCl = 10 to 29 mL/min: 100 mg</p> <p>Dose adjustment in pediatric patients 2 to 12 years of age with CrCl = 10 to 29 mL/min: 2 mg/kg</p> <p>HD: Administer after dialysis</p>
<p>Relenza (zanamivir)</p>	<p>Inhalation powder (in blisters)</p>	<p>Oral inhalation via Diskhaler device</p>	<p>Once daily or twice daily, depending on the indication</p> <p><u>Treatment (≥ 7 years)</u>: 10 mg twice daily for 5 days</p> <p><u>Prophylaxis in household setting (≥ 5 years)</u>: 10 mg once daily for 10 days</p>	<p>The 10-mg dose is provided by 2 inhalations (one 5-mg blister per inhalation).</p> <p>Patients scheduled to use an inhaled bronchodilator at the same time as Relenza should use their bronchodilator before taking Relenza.</p>

Drug	Available Formulations	Route	Usual Recommended Frequency	Comments
			<u>Prophylaxis in community outbreak (adults and adolescents):</u> 10 mg once daily for 28 days	<p>If Relenza is prescribed for children, it should be used only under adult supervision and instruction, and the supervising adult should first be instructed by a healthcare professional.</p> <p>Due to the low systemic bioavailability of Relenza following oral inhalation, no dosage adjustments are necessary for patients with renal Impairment; however, the potential for drug accumulation should be considered.</p>
Tamiflu (oseltamivir)	Capsules, powder for oral suspension	Oral	<p>Once daily or twice daily, depending on the indication</p> <p>Patients ≥ 13 years <u>Treatment:</u> 75 mg twice daily for 5 days</p> <p><u>Prophylaxis:</u> 75 mg once daily for at least 10 days following close contact with an infected individual and up to 6 weeks during a community outbreak. In immunocompromised patients, may be continued for up to 12 weeks.</p> <p>Patients < 13 years <u>Treatment:</u></p> <ul style="list-style-type: none"> • 2 weeks to < 1 year: 3 mg/kg twice daily for 5 days • 1 to 12 years: 30 to 75 mg twice daily for 5 days; specific weight-based dosing recommendations as follows: <ul style="list-style-type: none"> ○ ≤ 15 kg: 30 mg twice daily ○ 15.1 kg to 23 kg: 45 mg twice daily ○ 23.1 kg to 40 kg: 60 mg twice daily ○ ≥ 40.1 kg: 75 mg twice daily <p><u>Prophylaxis:</u></p> <ul style="list-style-type: none"> • 1 to 12 years: 30 to 75 mg 	<p>Start treatment within 48 hours of symptom onset or close contact with the infected individual.</p> <p>Taking with food may enhance tolerability. In an emergency, a suspension can be made from capsules.</p> <p>Dosage adjustment is recommended for patients with a CrCl between 10 and 60 mL/minute and for patients with ESRD undergoing routine HD or CAPD.</p> <p>Not recommended for patients with ESRD not undergoing dialysis.</p> <p>No dosage adjustment for mild to moderate hepatic impairment.</p> <p>Safety not evaluated in patients with severe hepatic impairment.</p>

Drug	Available Formulations	Route	Usual Recommended Frequency	Comments
			once daily for 10 days; specific weight-based dosing recommendations as follows: <ul style="list-style-type: none"> ○ ≤ 15 kg: 30 mg once daily ○ 15.1 kg to 23 kg: 45 mg once daily ○ 23.1 kg to 40 kg: 60 mg once daily ○ ≥ 40.1 kg: 75 mg once daily ● During a community outbreak, can continue for up to 6 weeks (or up to 12 weeks in immunocompromised patients). 	

CAPD=continuous ambulatory peritoneal dialysis; CrCl =creatinine clearance; ESRD=end stage renal disease; HD=hemodialysis
 *See the current prescribing information for full details

CONCLUSION

- The first line of protection against influenza is vaccination. All individuals six months of age and older without contraindications should receive yearly influenza vaccination (*AAP 2017, Fiore et al 2011, Grohskopf et al 2017*).
- Antivirals are available for the prevention and treatment of influenza. Overall, the adamantanes and neuraminidase inhibitors have demonstrated safety and efficacy for their respective indications. However, amantadine and rimantadine are not currently recommended due to high rates of resistance in circulating influenza virus strains (*CDC 2018*).
- Relenza (zanamivir) and oseltamivir are both effective in preventing influenza but are not substitutes for annual vaccination. They are recommended as post-exposure chemoprophylaxis in patients with a high risk for influenza complications who are not sufficiently protected by vaccination (*Fiore et al 2011, CDC 2018, Harper et al 2009, Panel on Opportunistic Infections 2013*). Rapivab (peramivir) is not approved or recommended for influenza prophylaxis (*CDC 2018*).
- Rapivab (peramivir), Relenza (zanamivir), and oseltamivir effectively treat influenza by reducing the duration of fever and illness. Initiation of treatment is recommended as soon as possible for patients with suspected influenza who are hospitalized, severely ill, or at high risk for influenza complications (*Fiore et al 2011, AAP 2017, CDC 2017, CDC 2018, Harper et al 2009, Panel on Opportunistic Infections 2013*).
- Limited within-class comparisons prevent the recommendation of one neuraminidase inhibitor over another. Factors to consider when selecting an antiviral agent include the route of administration, seasonal and geographical susceptibility trends, and patient-specific factors such as age and compliance (*Takemoto et al 2013*).
- The most common adverse events with amantadine and rimantadine are **nausea, insomnia, dizziness**, headache, anorexia, dry mouth, and agitation. The adamantanes are associated with an increased risk for seizures.
- The most common adverse events with Relenza (zanamivir) and oseltamivir are headache, nausea, and vomiting. Diarrhea is the most common adverse event with Rapivab (peramivir). The neuraminidase inhibitors have a labeled warning for neuropsychiatric events such as delirium and abnormal behavior leading to injury.

REFERENCES

- Amantadine capsules [package insert], Sunrise, FL: Cipla USA, Inc.; May 2017.
- Amantadine oral solution [package insert], Farmville, NC: CMP Pharm, Inc.; January 2015.
- Amantadine tablets [package insert], Parsippany, NJ: Actavis Pharma, Inc.; January 2017.
- American Academy of Pediatrics Committee on Infectious Diseases. Recommendations for prevention and control of influenza in children, 2017-2018. *Pediatrics*. 2017;140(4):e20172550.
- Anekthananon T, Pukritayakamee S, Ratanasuwon W, et al. Oseltamivir and inhaled zanamivir as influenza prophylaxis in Thai health workers: a randomized, double-blind, placebo-controlled safety trial over 16 weeks. *J Antimicrob Chemother*. 2013;68(3): 697-707.

- Aoki FY, Macleod MD, Paggiaro P, et al. Early administration of oral oseltamivir increases the benefits of influenza treatment: IMPACT trial. *J Antimicrob Chemother.* 2003;51(1):123-9.
- Birnkrant D, Cox E. The Emergency Use Authorization of Peramivir for Treatment of 2009 H1N1 Influenza. *N Engl J Med.* 2009;361(23):2204-2207.
- Bryson YJ, Monahan C, Pollack M, Shields WD. A prospective double-blind study of side effects associated with the administration of amantadine for influenza A virus prophylaxis. *J Infect Dis.* 1980;141(5):543-7.
- Centers for Disease Control and Prevention. Clinical Signs and Symptoms of Influenza. Updated May 26, 2016[a]. <https://www.cdc.gov/flu/professionals/acip/clinical.htm>. Accessed May 8, 2018.
- Centers for Disease Control and Prevention. Flu Symptoms & Complications. Updated May 23, 2016[b]. <https://www.cdc.gov/flu/about/disease/complications.htm>. Accessed May 8, 2018.
- Centers for Disease Control and Prevention. Influenza Antiviral Medications: Summary for Clinicians. Updated February 23, 2018. <http://www.cdc.gov/flu/professionals/antivirals/summary-clinicians.htm>. Accessed May 8, 2018.
- Centers for Disease Control and Prevention. CDC Health Advisory: Seasonal influenza A (H3N2) activity and antiviral treatment of patients with influenza. December 27, 2017. <https://emergency.cdc.gov/han/han00409.asp>. Accessed May 8, 2018.
- Centers for Disease Control and Prevention. Termination of Declaration of Emergency Justifying Emergency Use Authorization (EUA) of Certain Antiviral Drugs – Zanamivir, Oseltamivir Phosphate and Peramivir. June 21, 2010[a]. <http://www.cdc.gov/H1N1flu/EUA/pdf/Antivirals.pdf>. Accessed May 8, 2018.
- Centers for Disease Control and Prevention. Update: influenza activity - United States, 2009-10 season. *MMWR.* 2010[b]; 59(29):901-8.
- Chik KW, Li CK, Chan PK, et al. Oseltamivir prophylaxis during the influenza season in a pediatric cancer centre: prospective observational study. *Hong Kong Med J.* 2004;10(2):103-6.
- Cooper NJ, Sutton AJ, Abrams KR, et al. Effectiveness of neuraminidase inhibitors in treatment and prevention of influenza A and B: systematic review and meta-analyses of randomized controlled trials. *BMJ.* 2003;326(7401):1235.
- Crawford SA, Clover RD, Abell TD, et al. Rimantadine prophylaxis in children: a follow-up study. *Pediatr Infect Dis J.* 1988;7(6):379-83.
- De Jong MD, Ison MG, Monto AS, et al. Evaluation of intravenous peramivir for treatment of influenza in hospitalized patients. *Clin Inf Dis.* 2014;59(15 December):e172-e185.
- Dolin R, Reichman RC, Madore HP, et al. A controlled trial of amantadine and rimantadine in the prophylaxis of influenza A infection. *N Engl J Med.* 1982;307(10):580-4.
- Doll MK, Winters N, Boikos C, et al. Safety and effectiveness of neuraminidase inhibitors for influenza treatment, prophylaxis, and outbreak control: a systematic review of systematic reviews and/or meta-analyses. *J Antimicrob Chemother.* 2017;72:2990-3007.
- Drugs@FDA: FDA approved drug products. 2018. Food and Drug Administration Web site. <https://www.accessdata.fda.gov/scripts/cder/daf/>. Accessed May 8, 2018.
- Fiore AE, Frye A, Shay D, et al. Antiviral agents for the treatment and chemoprophylaxis of influenza: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm Rep.* 2011;60(1):1-24.
- Flumadine [package insert], Detroit, MI: Caraco Pharmaceutical Laboratories, Ltd.; April 2010.
- Food and Drug Administration. Center for Drug Evaluation and Research. New Drug Application – 206426Orig1s000. Peramivir Medical Review. Food and Drug Administration. August 22, 2014. http://www.accessdata.fda.gov/drugsatfda_docs/nda/2014/206426Orig1s000MedR.pdf. Accessed May 8, 2018.
- Fry AM, Goswami D, Nahar K, et al. Efficacy of oseltamivir treatment started within 5 days of symptom onset to reduce influenza illness duration and virus shedding in an urban setting in Bangladesh: a randomized placebo-controlled trial. *Lancet Infect Dis.* 2014;14(2):109-118.
- Grohskopf LA, Sokolow LZ, Broder KR, et al. Prevention and Control of Influenza with Vaccines: Recommendations of the Advisory Committee on Immunization Practices (ACIP) – United States, 2017-18 Influenza Season. August 25, 2017. *MMWR.* 2017; 66(2):1–20. <https://www.cdc.gov/mmwr/volumes/66/rr/rr6602a1.htm>. Accessed May 8, 2018.
- Hall CB, Dolin R, Gala CL, et al. Children with influenza A infection: treatment with rimantadine. *Pediatrics.* 1987;80(2):275-82.
- Halloran ME, Hayden FG, Yang Y, et al. Antiviral effects on influenza viral transmission and pathogenicity: observations from household-based trials. *Am J Epidemiol.* 2007;165(2):212-21.
- Harper SA, Bradley JS, Englund JA, et al. Seasonal influenza in adults and children—diagnosis, treatment, chemoprophylaxis, and institutional outbreak management: clinical practice guidelines of the Infectious Diseases Society of America. *Clin Infect Dis.* 2009;48(8):1003-32.
- Hayden FG, Atmar RL, Schilling M, et al. Use of the selective oral neuraminidase inhibitor oseltamivir to prevent influenza. *N Engl J of Med.* 1999;341(18):1336-43.
- Hayden FG, Belshe R, Villanueva C, et al. Management of influenza in households: a prospective, randomized comparison of oseltamivir treatment with or without post exposure prophylaxis. *J Infect Dis.* 2004;189(3):440-9.
- Hayden FG, Belshe RB, Clover RD, et al. Emergence and apparent transmission of rimantadine-resistant influenza A virus in families. *N Engl J Med.* 1989;321(25):1696-702.
- Hayden FG, Gubareva LV, Monto AS, et al. Inhaled zanamivir for the prevention of influenza in families. Zanamivir Family Study Group. *N Engl J Med.* 2000;343(18):1282-9.
- Hayden FG, Osterhaus AD, Treanor JJ, et al. Efficacy and safety of the neuraminidase inhibitor zanamivir in the treatment of influenza virus infections. GG167 Influenza Study Group. *N Engl J Med.* 1997;337(13):874-80.
- Hedrick JA, Barzilai A, Behre U, et al. Zanamivir for treatment of symptomatic influenza A and B infection in children five to twelve years of age: a randomized controlled trial. *Pediatr Infect Dis J.* 2000;19(5):410-7.
- Hiba V, Chowdhury M, Levi-Vinograd I, et al. Benefit of early treatment with oseltamivir in hospitalized patients with documented 2009 influenza A (H1N1): retrospective cohort study. *J Antimicrob Chemother.* 2011;66:1150-5.
- Ison MG, Fraiz J, Heller B, et al. Intravenous peramivir for treatment of influenza in hospitalized patients. *Antivir Ther.* 2014;19(4):349-61.
- Ison MG, Hui DS, Clezy K, et al. A clinical trial of intravenous peramivir compared with oral oseltamivir for the treatment of seasonal influenza in hospitalized patients. *Antivir Ther.* 2013;19:651-661.
- Jackson RJ, Cooper KL, Tappenden P, et al. Oseltamivir, zanamivir and amantadine in the prevention of influenza: a systematic review. *J Infect.* 2011;62:14-25.

- Jefferson T, Demicheli V, Di Pietrantonj C, et al. Amantadine and rimantadine for influenza A in adults. *Cochrane Database Syst Rev*. 2006[a];(2):CD001169.
- Jefferson T, Demicheli V, Rivetti D, et al. Antivirals for influenza in healthy adults: systematic review. *Lancet*. 2006[b];367(9507):303-13.
- Jefferson T, Jones MA, Doshi P, et al. Neuraminidase inhibitors for preventing and treating influenza in healthy adults and children (Review). *Cochrane Database Syst Rev*. 2014;4: CD008965.
- Kaiser L, Wat C, Mills T, et al. Impact of oseltamivir treatment on influenza-related lower respiratory tract complications and hospitalizations. *Arch Intern Med*. 2003;163(14):1667-72.
- Kawai N, Ikematsu H, Iwaki N, et al. A comparison of the effectiveness of oseltamivir for the treatment of influenza A and influenza B: a Japanese multicenter study of the 2003-2004 and 2004-2005 influenza seasons. *Clin Infect Dis*. 2006;43(4):439-44.
- Kawai N, Ikematsu H, Iwaki N, et al. A comparison of the effectiveness of zanamivir and oseltamivir for the treatment of influenza A and B. *J Infect*. 2008;56(1):51-7.
- Kawai N, Ikematsu H, Iwaki N, et al. Factors influencing the effectiveness of oseltamivir and amantadine for the treatment of influenza: a multicenter study from Japan of the 2002-2003 influenza season. *Clin Infect Dis*. 2005;40(9):1309-16.
- Kohno S, Kida H, Mizuguchi M, et al. Efficacy and safety of intravenous peramivir for treatment of seasonal influenza virus infection. *Antimicrob Agents Chemother*. 2010;54(11):4568-4574.
- Kohno S, Yen MY, Cheong HJ, et al. Phase III randomized, double-blind study comparing single-dose intravenous peramivir with oral oseltamivir in patients with seasonal influenza virus infection. *Antimicrob Agents Chemother*. 2011;55(11):5267-76.
- Lee J, Park J, Jwa H, Kim YH. Comparison of efficacy of intravenous peramivir and oral oseltamivir for the treatment of influenza: systematic review and meta-analysis. *Yonsei Med J*. 2017;54(4):778-785.
- Li TC, Chan MC, Lee N. Clinical implications of antiviral resistance in influenza. *Viruses*. 2015;7(9):4929-44.
- Lin JT, Yu XZ, Cui DJ, et al. A multicentre, randomized, controlled trial of oseltamivir in the treatment of influenza in a high-risk Chinese population. *Curr Med Res Opin*. 2006;22(1):75-82.
- MIST Study Group. Randomized trial of efficacy and safety of inhaled zanamivir in treatment of influenza A and B virus infections. The MIST (Management of Influenza in the Southern Hemisphere Trialists) Study Group. *Lancet*. 1998;352(9144):1877-81.
- Monto A, Ohmit SE, Hornbuckle K, et al. Safety and efficacy of long-term use of rimantadine for prophylaxis of type A influenza in nursing homes. *Antimicrob Agents Chemother*. 1995;39(10):2224-8.
- Monto AS, Fleming DM, Henry D, et al. Efficacy and safety of the neuraminidase inhibitor zanamivir in the treatment of influenza A and B virus infections. *J Infect Dis*. 1999[a];180(2):254-61.
- Monto AS, Pichichero ME, Blanckenberg SJ, et al. Zanamivir prophylaxis: an effective strategy for the prevention of influenza types A and B within households. *J Infect Dis*. 2002;186(11):1582-8.
- Monto AS, Robinson DP, Herlocher ML, et al. Zanamivir in the prevention of influenza among healthy adults: a randomized controlled trial. *JAMA*. 1999[b];282(1):31-5.
- Nicholson KG, Aoki FY, Osterhaus AD, et al. Efficacy and safety of oseltamivir in treatment of influenza: a randomized controlled trial. *Lancet*. 2000;355(9218):1845-50.
- Orange Book: Approved drug products with therapeutic equivalence evaluations. 2018. Food and Drug Administration Web site. <https://www.accessdata.fda.gov/scripts/cder/ob/default.cfm>. Accessed May 8, 2018.
- Panel on Opportunistic Infections in HIV-Exposed and HIV-Infected Children. Guidelines for the prevention and treatment of opportunistic infections in HIV-exposed and HIV-infected children. Department of Health and Human Services. Updated November 6, 2013. https://aidsinfo.nih.gov/contentfiles/lvguidelines/oi_guidelines_pediatrics.pdf. Accessed May 8, 2018.
- Peters PH Jr, Gravenstein S, Norwood P, et al. Long-term use of oseltamivir for the prophylaxis of influenza in a vaccinated frail older population. *J Am Geriatr Soc*. 2001;49(8):1025-31.
- Rapivab [package insert], Durham, NC: BioCryst Pharmaceuticals, Inc.; April 2018.
- Relenza [package insert], Research Triangle Park, NC: GlaxoSmithKline LLC; August 2016.
- Reuman PD, Bernstein DI, Keefer MC, et al. Efficacy and safety of low dosage amantadine hydrochloride as prophylaxis for influenza A. *Antiviral Res*. 1989;11(1):27-40.
- Singh S, Barghoorn J, Bagdonas A, et al. Clinical benefits with oseltamivir in treating influenza in adult populations: results of a pooled and subgroup analysis. *Clin Drug Investig*. 2003;23(9):561-9.
- Sugaya N, Kohno S, Ishibashi T, et al. Efficacy, safety, and pharmacokinetics of intravenous peramivir in children with 2009 pandemic H1N1 influenza A virus infection. *Antimicrob Agents Chemother*. 2012;56(1):369-377.
- Takemoto Y, Asai T, Ikezoe I, et al. Clinical effects of oseltamivir, zanamivir, laninamivir and peramivir on seasonal influenza infection in outpatients in Japan during the winter of 2012-2013. *Chemotherapy*. 2013;59(5):373-8.
- Tamiflu [package insert], South San Francisco, CA: Genentech, Inc.; April 2018.
- Treanor JJ, Hayden FG, Vrooman PS, et al. Efficacy and safety of the oral neuraminidase inhibitor oseltamivir in treating acute influenza. *JAMA*. 2000;283(8):1016-24.
- Turner D, Wailoo A, Nicholson K, et al. Systematic review and economic decision modeling for the prevention and treatment of influenza A and B. *Health Technol Assess*. 2003;7(35):iii-iv, xi-xiii, 1-170.
- Wang K, Shun-Shin M, Gill P, et al. Neuraminidase inhibitors for preventing and treating influenza in children. *Cochrane Database Syst Rev*. 2012;4:CD002744.
- Welliver R, Monto A, Carewicz O, et al. Effectiveness of oseltamivir in preventing influenza in household contacts. *JAMA*. 2001;285(6):748-54.
- Whitley RJ, Hayden FG, Reisinger KS, et al. Oral oseltamivir treatment of influenza in children. *Pediatr Infect Dis J*. 2001;20(2):127-33.

Publication Date: July 2, 2018