

Therapeutic Class Overview

Neuropathic Pain and Fibromyalgia Agents

INTRODUCTION

- Neuropathic pain is commonly described by patients as burning or electrical in nature and results from injury or damage to the nervous system (*Herndon et al 2017*). Management of neuropathic pain may prove challenging due to unpredictable patient response to drug therapy (*Attal et al 2010*).
- Fibromyalgia is characterized by chronic musculoskeletal pain with unknown etiology and pathophysiology. Patients typically complain of widespread musculoskeletal pain, fatigue, cognitive disturbance, psychiatric symptoms, and multiple somatic symptoms (*Goldenberg 2016[a]*). Fibromyalgia is often difficult to treat and requires a multidisciplinary, individualized treatment program (*Goldenberg 2016[b]*).
- This review focuses on medications that are approved by the Food and Drug Administration (FDA) for the treatment of fibromyalgia, neuropathic pain, and/or post-herpetic neuralgia (PHN). The products in this review include Cymbalta (duloxetine), Gralise (gabapentin ER), Horizant (gabapentin enacarbil ER), Lidoderm (lidocaine 5% patch), Lyrica (pregabalin), Lyrica CR (pregabalin ER), Neurontin (gabapentin), Nucynta ER (tapentadol ER), Qutenza (capsaicin), and Savella (milnacipran). These agents represent a variety of pharmacologic classes, including anticonvulsants, serotonin-norepinephrine reuptake inhibitors (SNRIs), extended-release (ER) opioids, and topical analgesics. As such, these agents hold additional FDA-approved indications that are outlined in Table 2; however, clinical information included within this review will not address the use of these agents for these additional indications (*Prescribing information: Cymbalta 2017, Gralise 2015, Horizant 2016, Lidoderm 2015, Lyrica 2017, Lyrica CR 2017, Neurontin 2017, Nucynta ER 2017, Qutenza 2013, Savella 2017*).
 - In February 2018, the FDA approved ZTlido (lidocaine 1.8% topical system) for the relief of pain associated with PHN. The product is not yet available commercially.
 - One ZTlido 1.8% topical system provides equivalent lidocaine exposure to one Lidoderm 5% patch.
- Medispan classes: Anticonvulsants - Misc.; Fibromyalgia Agents; Local Anesthetics – Topical; Opioid Agonists; Postherpetic Neuralgia (PHN) Agents; Restless Leg Syndrome (RLS) Agents; Serotonin-Norepinephrine Reuptake Inhibitors (SNRIs)

Diabetic Neuropathy

- Approximately 50% of patients with diabetes will eventually develop neuropathy. The high rate of diabetic neuropathy results in substantial patient morbidity, which includes recurrent lower extremity infections, ulcerations, and subsequent amputations (*Feldman et al 2015[a]*).
- The condition is categorized into distinct syndromes based on the neurologic distribution, although syndromes may overlap in some patients. The most frequently encountered diabetic neuropathies include distal symmetric polyneuropathy, autonomic neuropathy, polyradiculopathies, and mononeuropathies (*Feldman et al 2015[a]*).
- The 3 main components to the management of diabetic neuropathy are glycemic control, foot care, and pain management (*Feldman et al 2017[b]*).
 - Optimal glucose control is important for the prevention of diabetic neuropathy. Clinical trial evidence demonstrates that rigorous blood glucose control in patients with type 1 diabetes reduces the occurrence of diabetic neuropathy. In contrast, the role of glycemic control in established diabetic neuropathy is uncertain. Limited evidence suggests that neuropathic symptoms may improve with intensive antidiabetic therapy (*Feldman et al 2017[b]*).
 - Patients with diabetes should be counseled on the importance of daily foot care, including the inspection of feet for the presence of dry or cracking skin, fissures, and plantar callus formation. Regular foot examinations by a healthcare provider are also important (*Feldman et al 2017[b]*).
 - A small proportion of patients with diabetic neuropathy will experience painful symptoms, and in some instances the condition is self-limited. When treatment is necessary, options include antidepressants, anticonvulsants, capsaicin cream, lidocaine patches, alpha-lipoic acid, isosorbide dinitrate topical spray, and transcutaneous electrical nerve stimulation (*Feldman et al 2017[b]*).

Fibromyalgia

- Fibromyalgia is a chronic functional illness marked by widespread musculoskeletal pain for which no alternative cause can be identified. Fibromyalgia patients often experience neuropsychological symptoms of fatigue, unrefreshing sleep, cognitive dysfunction, anxiety, and depression (*Clauw et al 2009*).
 - Patients with fibromyalgia have pain that is typically above and below the waist on both sides of the body and involves the axial skeleton (neck, back, or chest). The pain attributable to fibromyalgia is poorly localized, difficult to ignore, severe in its intensity, and associated with a reduced functional capacity (*Crofford 2015*).
- The prevalence of fibromyalgia in the general US population is estimated to be 2% (*Williams et al 2009*). It is more common in women than in men, with a ratio of approximately 9:1 (*Crofford 2015*).
- There is an increased prevalence of other syndromes associated with pain and fatigue, including chronic fatigue syndrome, temporomandibular disorder, chronic headaches, irritable bowel syndrome, interstitial cystitis/painful bladder syndrome, and other pelvic pain syndromes in fibromyalgia patients (*Clauw et al 2009, Crofford 2015*).

PHN

- PHN is generally defined as the persistence of the pain of herpes zoster for more than three months after resolution of the rash. It affects 10 to 15% of patients with herpes zoster, with incidence increasing with age. The duration of PHN is highly variable among individuals (*Dubinsky et al 2004*).
- PHN, as well as acute herpetic neuralgia, can be a severe condition associated with profound psychological dysfunction, including impaired sleep, decreased appetite, and decreased libido (*Bajwa et al 2017*).
- Prevention of PHN involves either treatment of acute herpes zoster infection or use of a vaccine (*Bajwa et al 2017*). Administration of antiviral agents within 72 hours of the onset of herpes zoster can reduce the intensity and duration of acute illness, and can also prevent PHN. This may also be achieved with the administration of the tricyclic antidepressant, amitriptyline (*Dubinsky et al 2004*).
- A number of treatment modalities have been evaluated in the management of PHN and include tricyclic antidepressants, anticonvulsants, opioids, capsaicin, topical lidocaine, intrathecal glucocorticoids, N-methyl-D-aspartate receptor antagonists, botulinum toxin, cryotherapy, and surgery (*Bajwa et al 2017*).

Table 1. Medications Included Within Class Review

Drug	Generic Availability
Cymbalta (duloxetine delayed-release)	✓
Gralise (gabapentin ER)*	-
Horizant (gabapentin enacarbil ER)*	-
Lidoderm (lidocaine transdermal patch)	✓
Lyrica (pregabalin)	-
Lyrica CR (pregabalin ER)	-
Neurontin (gabapentin)*	✓
Nucynta ER (tapentadol ER)	-
Qutenza (capsaicin transdermal patch)	-
Savella (milnacipran)	-
ZTlido (lidocaine topical system)	†

* Medication is not interchangeable with other gabapentin products because of differing pharmacokinetic profiles that affect the frequency of administration.

† ZTlido is not yet available commercially.

(*Drugs@FDA 2018, Orange Book: Approved Drug Products with Therapeutic Equivalence Evaluations 2018*)

INDICATIONS
Table 2. FDA-Approved Indications

Indication	Cymbalta (duloxetine)	Gralise (gabapentin ER)	Horizant (gabapentin enacarbil ER)	Lidoderm, ZTlido (lidocaine)	Lyrica (pregabalin)	Lyrica CR (pregabalin ER)	Neurontin (gabapentin)	Nucynta ER (tapentadol)	Qutenza (capsaicin)	Savella (milnacipran)
Adjunctive therapy for adult patients with partial onset seizures					✓					
Adjunctive therapy in the treatment of partial seizures with and without secondary generalization in patients > 3 years of age with epilepsy							✓			
Adjunctive therapy for patients 4 years of age and older with partial onset seizures					✓					
Management of chronic musculoskeletal pain	✓ †									
Management of fibromyalgia	✓				✓					✓
Management of neuropathic pain associated with diabetic peripheral neuropathy	✓				✓	✓		✓ §		
Management of neuropathic pain associated with spinal cord injury					✓					
Management of PHN		✓	✓		✓	✓	✓			
Relief of pain associated with PHN				✓					✓	
Moderate-to-severe primary restless legs syndrome			✓ †							
Treatment of generalized anxiety disorder	✓									
Treatment of major depressive disorder	✓									
Management of moderate to severe chronic pain in adults								✓ §		

† This has been established in studies of patients with chronic low back pain and chronic pain due to osteoarthritis.

‡ Gabapentin enacarbil is not indicated for patients who are required to sleep during the day and remain awake at night.

§ Indicated when a continuous, around-the-clock opioid analgesic is needed for an extended period of time. Medication is not for use as: an as-needed analgesic; for pain that is mild or not expected to persist for an extended period of time; for acute pain; or for postoperative pain, unless the patient is already receiving chronic opioid therapy prior to surgery, or if the postoperative pain is expected to be moderate to severe and persist for an extended period of time.

(Prescribing information: Cymbalta 2017, Gralise 2015, Horizant 2016, Lidoderm 2015, Lyrica 2018, Lyrica CR 2017, Neurontin 2017, Nucynta ER 2017, Qutenza 2013, Savella 2017, ZTlido 2018)

- Information on indications, mechanism of action, pharmacokinetics, dosing, and safety has been obtained from the prescribing information for the individual products, except where noted otherwise.

CLINICAL EFFICACY SUMMARY

Neuropathic Pain

- Pregabalin demonstrated significant improvements in pain relief, functional outcomes, and quality of life compared to placebo for the treatment of diabetic peripheral neuropathic pain. Commonly reported adverse events (AEs) in patients receiving pregabalin include dizziness, somnolence, infection, headache, dry mouth, weight gain, and peripheral edema (*Dworkin et al 2003, Freynhagen et al 2005, Guan et al 2011, Lesser et al 2004, Moon et al 2010, Rosenstock et al 2004, Roth et al 2010, Sabatowski et al 2004, Semel et al 2010, Sharma et al 2010, Skvarc et al 2010*).
- Tapentadol ER demonstrated superiority over placebo in alleviating pain and improving quality of life in patients with diabetic peripheral neuropathy. Tapentadol ER is associated with significant improvements in pain intensity scores, responder rates, and patient global impression of change (PGIC). Commonly reported AEs in patients receiving tapentadol ER include nausea, vomiting, and constipation (*Schwartz et al 2011*).
- Duloxetine demonstrated consistent superiority over placebo in alleviating pain, improving functional outcomes, and improving quality of life in patients with diabetic peripheral neuropathic pain. Specifically, duloxetine is associated with significant improvements in Brief Pain Inventory, Clinician and Patient Global Impression of Improvement and Severity, Short Form-36 Health Survey (SF-36), Pain-Related Sleep Interference, and Euro Quality of Life assessment (EQ-5D) scores. Commonly reported AEs in patients receiving duloxetine include nausea, somnolence, anorexia, and dysuria (*Armstrong et al 2007, Kajdasz et al 2007, Lunn et al 2014, Parsons et al 2016, Yan et al 2010*).
- Head-to-head trials among the neuropathic pain and fibromyalgia agents are rare. In a 52-week, open-label trial comparing duloxetine to routine care (gabapentin, amitriptyline, and venlafaxine) for the treatment of diabetic peripheral neuropathic pain, there were no significant differences observed between groups in EQ-5D questionnaire scores; however, results differed with regards to SF-36 subscale scores. In another trial, there were no significant between-group differences in SF-36 subscale scores; however, other subscale scores for physical functioning, bodily pain, mental health, and vitality favored duloxetine (*Raskin et al 2006, Wernicke et al 2007[b]*). A second head-to-head trial demonstrated duloxetine to be noninferior to pregabalin for the treatment of pain in patients with diabetic peripheral neuropathy who had an inadequate pain response to gabapentin (*Tanenberget al 2011*). A post-hoc analysis of study patients who were taking concomitant antidepressants and those who were not taking antidepressants found duloxetine may provide better pain reduction in those patients who were not taking concomitant antidepressants (*Tanenberget al 2014*). Another head-to-head trial found high-dose duloxetine or pregabalin monotherapy had no significant differences, as measured by Brief Pain Inventory Modified Short Form (BPI-MSF) average pain in comparison, with combination duloxetine and pregabalin therapy (*Tesfaye et al 2013*).
- Several large meta-analyses and systematic reviews have been conducted evaluating the neuropathic pain and fibromyalgia agents, which further support the safety and efficacy of these agents in FDA-approved indications (*Chou et al 2009, Edelsberg et al 2011, Lunn et al 2009, Lunn et al 2014, Meng et al 2014, Moore et al 2009, Moore et al 2014, Quilici et al 2009, Wernicke et al 2007[a]*). In a meta-analysis by Quilici et al, limited available clinical trial data suitable for indirect comparison demonstrated that duloxetine provides comparable efficacy and tolerability to that of gabapentin and pregabalin for the treatment of diabetic peripheral neuropathic pain (*Quilici et al 2009*).
- The efficacy of pregabalin in patients with neuropathic pain associated with spinal cord injury was established in 2 placebo-controlled trials, 1 of 12 weeks duration and the other of 16 weeks duration. Patients had neuropathic pain associated with spinal cord injury for at least 3 months or with relapses and remissions for at least 6 months. Patients were allowed to take opioids, non-opioid analgesics, antiepileptic drugs, muscle relaxants, and antidepressant drugs if doses were stable for 30 days prior to screening. Patients were also allowed to take acetaminophen and nonsteroidal anti-inflammatory drugs during the trial. In both trials, pregabalin (150 to 600 mg/day) significantly improved weekly pain scores, and increased the proportion of patients with at least a 30 or 50% reduction from baseline in pain score compared to placebo (*Lyrice prescribing information 2016, Siddall et al 2006, Vranken et al 2008*).

Fibromyalgia

- From the agents included in this review, the agents that have several randomized controlled trials (RCTs) and meta-analyses demonstrating their efficacy in the treatment of fibromyalgia include duloxetine, pregabalin, and milnacipran (*Arnold et al 2007, Arnold et al 2008, Arnold et al 2009, Clauw et al 2008, Crofford et al 2005, Hauser et al 2009[a], Hauser et al 2009[b], Hauser et al 2010, Hauser et al 2013, Lunn et al 2014, Mease et al 2009, Mease et al 2010, Russell et al 2008, Vitton et al 2004*).
 - A 2009 meta-analysis on the treatment of fibromyalgia syndrome with antidepressants found that antidepressants were associated with improved health-related quality of life. The largest effect size for pain reduction was seen with

the tricyclic antidepressant, amitriptyline, followed by monoamine oxidase inhibitors, moclobemide and pirlindole (medium effect size). Small effect sizes were observed with the selective serotonin reuptake inhibitors (SSRIs), fluoxetine and paroxetine, and the SNRIs, duloxetine and milnacipran. The authors concluded that short-term treatment with amitriptyline and duloxetine could be considered for fibromyalgia-associated pain and sleep disturbances (*Hauser et al 2009[a]*).

- In a meta-analysis of 5 RCTs, gabapentin and pregabalin reduced pain and improved sleep in patients with fibromyalgia. The pooled number-needed-to-treat to achieve $\geq 30\%$ reduction in pain was 8.5. Anxiety, depressed mood, and fatigue were not improved with gabapentin or pregabalin treatment (*Hauser et al 2009[b]*).
- Results from another 2010 meta-analysis noted that duloxetine, milnacipran, and pregabalin have short-term (up to 6-month) efficacy data. The authors concluded that the choice of medication may be dependent on the differences with regard to the occurrence of the key symptoms of fibromyalgia syndrome and to AEs specific to individual drug (*Hauser et al 2010*).
- A systematic review and meta-analysis of 10 randomized trials involving 6038 patients concluded that duloxetine and milnacipran provided a small incremental benefit over placebo in pain reduction, and that the dropout rates for duloxetine and milnacipran due to AEs were higher than placebo (*Hauser et al 2013*).
- A systematic review of 6 randomized trials involving 2249 patients concluded that for the treatment of fibromyalgia, duloxetine 60 and 120 mg/day are effective with a similar magnitude of effect (low quality evidence). The effect in fibromyalgia may be achieved through a greater improvement in mental symptoms than somatic physical pain (*Lunn et al 2014*).
- A 2016 network meta-analysis of 9 RCTs (N = 5140) indirectly compared duloxetine, pregabalin, and milnacipran in the treatment of fibromyalgia. The probability of achieving $> 30\%$ improvement in pain scores was numerically highest with duloxetine 60 mg, followed by pregabalin 300 mg, milnacipran 100 mg, and milnacipran 200 mg. While the aforementioned treatment groups each demonstrated superiority over placebo, differences between active treatments did not achieve statistical significance (*Lee and Song 2016*).

PHN

- In patients with PHN, treatment with lidocaine 5% resulted in significant pain relief compared to placebo (*Galer et al 1999, Galer et al 2002, Meier et al 2003*). In addition, treatment with lidocaine 5% was associated with higher rates of patient preference, less use of rescue medication, and decreases in allodynia and neuropathic symptoms compared to placebo (*Galer et al 1999, Meier et al 2003*). An open-label trial evaluating lidocaine 5% for the management of PHN supports the findings of placebo-controlled trials (*Katz et al 2002*).
- Lidocaine 1.8% was approved via the 505(b)(2) pathway with no new efficacy trials. However, in a single-dose, crossover study conducted in 53 healthy volunteers, lidocaine 1.8% topical system demonstrated equivalent exposure (AUC) and peak concentration (C_{max}) of lidocaine to lidocaine 5% patch. In addition, based on a clinical study in 54 subjects, 47 subjects (87%) had adherence scores of 0 ($\geq 90\%$ adhered) for all evaluations performed every 3 hours during the 12 hours of lidocaine 1.8% administration, 7 subjects (13%) had adherence scores of 1 ($\geq 75\%$ to $< 90\%$ adhered) for at least 1 evaluation, and no subjects had scores of 2 or greater ($< 75\%$ adhered) (*ZTlido prescribing information 2018*).
- In patients with PHN, treatment with capsaicin resulted in significant pain relief compared to low dose capsaicin 0.04% (*Backonja et al 2008, Derry et al 2017, Irving et al 2012*). Treatment with capsaicin was associated with improvement in PGIC, reduction in numeric pain rating scale (NPRS) scores, and reduction in neuropathic symptoms compared to low-dose capsaicin for up to 12 weeks of treatment (*Backonja et al 2008, Derry et al 2017, Irving et al 2012*). The long-term tolerability and safety of capsaicin was also demonstrated in a 52-week study, which found that repeat treatment with capsaicin (30 and 60 minutes) in addition to the standard of care therapies (antidepressants, antiepileptics, and/or opioids) was well tolerated with no negative functional or neurological effects when compared to standard of care therapies alone (*Vinik et al 2016*).
- Gabapentin also demonstrated superiority over placebo in alleviating pain, improving functional outcomes, and improving quality of life in patients with PHN. Treatment with gabapentin significantly improved average daily pain and sleep, Short form-McGill Pain Questionnaire (SF-MPQ), Patient and Clinician Global Impression of Change, SF-36, and Profile of Mood States (POMS) scores in RCTs. Commonly reported AEs in patients receiving gabapentin included somnolence, drowsiness, dizziness, ataxia, peripheral edema, and infection (*Rice et al 2001, Rowbotham et al 1998*). In a trial comparing placebo, gabapentin monotherapy, morphine sustained-release monotherapy, and gabapentin and morphine sustained-release combination therapy, combination therapy achieved better analgesia at lower doses of each

agent compared to monotherapy with either agent in patients with PHN. Combination therapy was most commonly associated with constipation, sedation, and dry mouth (*Gilron et al 2005*). Within these clinical trials, doses of gabapentin of up to 3,600 mg/day were evaluated (*Gilron et al 2005, Rice et al 2001, Rowbotham et al 1998*).

- In 2 placebo-controlled trials, gabapentin ER achieved significant improvements in average daily pain and sleep interference scores (*Irving et al 2009, Wallace et al 2010*). In one of these trials, a larger proportion of patients receiving gabapentin ER reported $\geq 50\%$ reduction from baseline in average daily pain scores compared to placebo (*Irving et al 2009*). In general, treatment with gabapentin ER was well tolerated; dizziness, headache, somnolence, and peripheral edema were the most commonly reported AEs (*Irving et al 2009, Wallace et al 2010*). Another placebo-controlled trial concluded that gabapentin ER may be particularly effective in patients with PHN presenting with sharp, dull, sensitive, or itchy pain (*Jensen et al 2009*). Within these clinical trials, doses of gabapentin ER of up to 1,800 mg/day were evaluated (*Irving et al 2009, Jensen et al 2009, Wallace et al 2010*).
- The efficacy of gabapentin enacarbil ER (1200, 2400, and 3600 mg/day) was established in a randomized, placebo-controlled, 12-week trial in adult patients with a documented medical diagnosis of PHN for ≥ 3 months (N = 371) and significant pain, as demonstrated by a minimum baseline 24-hour average Pain Intensity Numerical Rating Scale score ≥ 4 on the 11-point scale. Treatment with gabapentin enacarbil ER significantly improved the mean pain score and increased the proportion of patients with $\geq 50\%$ reduction in pain score from baseline at all doses evaluated. A benefit over placebo was observed for all 3 doses of gabapentin enacarbil ER as early as Week 1 and was maintained at Week 12. Additional benefit of using doses of gabapentin enacarbil ER > 1200 mg/day was not demonstrated (*Zhang et al 2013*). Results of a second, published, placebo-controlled trial confirms these findings. Reported AEs were similar to those of gabapentin and gabapentin ER (ie, dizziness, headache, and nausea) (*Backonja et al 2011*).
- A meta-analysis of 7 trials evaluating gabapentin, gabapentin enacarbil ER, and gabapentin ER was conducted to determine the efficacy and safety of all gabapentin formulations for management of PHN. Although gabapentin was found to be superior to placebo in terms of pain reduction, global impression of change, and sleep quality, patients taking gabapentin were significantly more likely to experience AEs such as dizziness, somnolence, peripheral edema, ataxia, and diarrhea (*Meng et al 2014*).
- Pregabalin demonstrated consistent superiority over placebo in alleviating diabetic peripheral neuropathic pain and PHN-related pain. Two noncomparative, open-label trials evaluating pregabalin for the management of PHN supports the findings of placebo-controlled trials (*Ogawa et al 2010, Xochilcal-Morales et al 2010*). In one of these noncomparative trials, long-term treatment of PHN with pregabalin (52 weeks) was found to be safe and effective (*Ogawa et al 2010*). Patients with PHN who were transitioned to pregabalin from gabapentin demonstrated no significant difference in pain scores, based on a visual analog scale, with pregabalin compared to gabapentin. However, in a subset of patients who required an increase in the dosage of pregabalin to improve the analgesic effect after the transition, significant improvement in pain scores was observed (*Ifuku et al 2011*).
- Support for efficacy of pregabalin ER in PHN and diabetic peripheral neuropathy was based on the efficacy of Lyrica in these indications and 1 clinical trial in PHN (*Lyrica CR prescribing information 2017*). In this trial, pregabalin ER demonstrated a significantly longer time to loss of therapeutic response compared with placebo over a 13-week randomized withdrawal phase in a phase 3, double-blind, randomized trial (*Huffman et al 2017*).

CLINICAL GUIDELINES

Diabetic Neuropathy

- The 2011 American Academy of Neurology (AAN) guidelines, which were reaffirmed in 2016, recommend the following:
 - If clinically appropriate, pregabalin should be offered for treatment. Gabapentin and sodium valproate are other anticonvulsants that should be considered for treatment (*Bril et al 2011*).
 - Amitriptyline, venlafaxine, and duloxetine should be considered for treatment; there is insufficient evidence available to recommend one of these agents over another. Combination therapy with venlafaxine and gabapentin may be utilized for a better response.
 - The opioids, dextromethorphan, morphine sulfate, tramadol, and oxycodone should be considered for treatment; there is insufficient evidence available to recommend one of these agents over another.
 - With regards to other pharmacologic options, capsaicin and isosorbide dinitrate spray should be considered for treatment, while lidocaine patch may be considered.
- The 2018 American Diabetes Association (ADA) guideline acknowledges the lack of quality of life outcomes and recommends that treatment decisions follow a trial-and-error approach (*ADA 2018*).

- Pregabalin, duloxetine, and tapentadol ER have been approved for relief of diabetic peripheral neuropathy; however, none of these agents affords complete relief, even when used in combination.
- Either pregabalin or duloxetine is recommended as initial pharmacologic therapy for neuropathic pain in diabetes. The use of tapentadol ER is generally not recommended as a first or second-line therapy due to safety concerns such as high-risk for addiction, and the evidence for its use is considered weaker.
- Venlafaxine, amitriptyline, gabapentin, carbamazepine, tramadol, and topical capsaicin may also be effective and could be considered for treatment of painful diabetic peripheral neuropathy.
- In general, other published guidelines support recommendations from the AAN and ADA concerning the use of the neuropathic pain and fibromyalgia agents in the management of diabetic neuropathy (*Dworkin et al 2007, Handelsman et al 2015, Pop-Busui et al 2017*).

PHN

- According to the 2004 AAN guidelines, first-line therapies for the management of PHN include tricyclic antidepressants, gabapentin, pregabalin, opioids, and topical lidocaine. The use of these therapies for long-term management remains uncertain (*Dubinsky et al 2004*).
- In general, other published guidelines support recommendations from the AAN for the use of the neuropathic pain and fibromyalgia agents in the management of PHN (*Attal et al 2010, Dworkin et al 2007*).

Fibromyalgia

- According to the evidence-based recommendations for the management of fibromyalgia syndrome from the European League Against Rheumatism, non-pharmacologic interventions should be considered first-line therapy for the management of fibromyalgia symptoms. Pharmacologic therapy should only be initiated if there is a lack of effect with non-pharmacologic therapies, and should be tailored to meet the patient's needs. Recommended pharmacologic agents include low-dose amitriptyline, cyclobenzaprine, duloxetine, milnacipran, pregabalin, and tramadol (*Macfarlane 2017*).
- According to the 2012 Canadian Guidelines for the diagnosis and management of fibromyalgia syndrome, all classes of antidepressants are options for treatment of pain and other symptoms of fibromyalgia. Anticonvulsants are also options, though the guideline does not recommend specific agents (*Fitzcharles et al 2013*).

SAFETY SUMMARY

- Duloxetine and milnacipran carry a boxed warning for clinical worsening, suicidality, and unusual changes in behavior. There is an increased risk of suicidal thinking and behavior in children, adolescents, and young adults taking antidepressants. All SNRIs are not approved for use in pediatric populations. All patients being treated with antidepressants for any indication should be monitored appropriately and observed closely, especially during the initial few months of a course of drug therapy and following changes in dosage.
- Tapentadol ER has a boxed warning for the potential for abuse, life-threatening respiratory depression, accidental exposure, risk of neonatal opioid withdrawal syndrome with prolonged use, and interactions with alcohol, benzodiazepines, or other central nervous system depressants that can cause profound sedation, respiratory depression, coma, and death.
- Gabapentin, pregabalin, and pregabalin ER carry warnings regarding the risk of anaphylaxis and/or angioedema after the first dose or during therapy.
- Topical lidocaine products have a warning for excessive dosing/overexposure, increased absorption on non-intact skin, risk of overexposure with external heat sources, and hypersensitivity reactions.
- The following key contraindications are included in the prescribing information:
 - Concomitant use or use within the last 14 days of monoamine oxidase inhibitors is contraindicated with duloxetine, milnacipran, and tapentadol ER.
 - Tapentadol ER is contraindicated in significant respiratory depression, acute or severe bronchial asthma, or hypercarbia in an unmonitored setting or in the absence of resuscitative equipment, and in known or suspected paralytic ileus.
- The FDA requires a Risk Evaluation and Mitigation Strategy (REMS) program for ER and long-acting opioid analgesics, including tapentadol ER, to assure safe use of these medications.
- The following monitoring parameters are recommended with treatment:
 - Monitor for clinical worsening of depression, suicidality, or unusual changes in behavior with duloxetine, milnacipran, gabapentin ER, gabapentin enacarbil ER, pregabalin, pregabalin ER, and gabapentin treatment.

- Patients receiving tapentadol ER, duloxetine, or milnacipran should be monitored for signs of serotonin syndrome when used concurrently with other serotonergic agents (eg, SSRIs, SNRIs, tricyclic antidepressants, triptans, fentanyl, lithium, tramadol, tryptophan, buspirone, amphetamines, and St. John's Wort). Tapentadol ER, duloxetine or milnacipran should not be used with drugs that impair metabolism of serotonin (eg, monoamine oxidase inhibitors [MAOIs], linezolid, and methylene blue).
- Monitor for signs of misuse, abuse, and addiction during tapentadol ER therapy. Patients should also be closely monitored for 72 hours after initiating tapentadol ER treatment and monitored throughout treatment due to an increased risk of respiratory depression.
- Patients receiving tapentadol ER, duloxetine, capsaicin, or milnacipran should have their blood pressure monitored prior to initiating treatment and periodically throughout treatment.
- Monitor for worsened seizure control in patients with a history of seizure disorder with the treatment of tapentadol ER, duloxetine, or milnacipran.
- Patients receiving tapentadol ER should be monitored for signs and symptoms of worsening biliary tract disease, including acute pancreatitis.
- In general, oral neuropathic pain and fibromyalgia agents are commonly associated with central nervous system-related AEs (eg, dizziness, drowsiness, somnolence). Peripheral edema and weight gain may also occur with use of these agents.

DOSING AND ADMINISTRATION

Table 3. Dosing and Administration

Drug	Available Formulations	Route	Usual Recommended Frequency	Comments
Cymbalta (duloxetine delayed-release)	Capsule	Oral	Once daily	Not recommended in ESRD, severe renal impairment (CrCl < 30 mL/min), or hepatic insufficiency
Gralise (gabapentin ER)	Tablet	Oral	Once daily	Administer with evening meal Reduce dose in CrCl of 30 to 60 mL/min; not recommended in CrCl < 30 mL/min or hemodialysis
Horizant (gabapentin enacarbil ER)	Tablet	Oral	Twice daily	Administer with food Reduce dose in CrCl < 60 mL/min or hemodialysis
Lidoderm, ZTlido (lidocaine)	Patch, topical system	Transdermal	Once daily	Apply for up to 12 hours within a 24-hour period Caution in patients with severe hepatic disease
Lyrica (pregabalin)	Capsule, oral solution	Oral	2 or 3 times daily	Schedule V controlled substance Reduce dose in CrCl < 60 mL/min
Lyrica CR (pregabalin ER)	Tablet	Oral	Once daily	Schedule V controlled substance Reduce dose in CrCl < 60 mL/min Administer after evening meal
Neurontin (gabapentin)	Capsule, oral solution, tablet	Oral	3 times daily	Reduce dose in CrCl < 60 mL/min
Nucynta ER (tapentadol ER)	Tablet	Oral	Twice daily	Schedule II controlled substance Do not use in severe renal impairment (CrCl < 30 mL/min) or severe hepatic impairment Reduce dose in moderate hepatic impairment
Qutenza (capsaicin)	Patch	Transdermal	60-minute application of up to 4 patches every 3 months	Only administered by physicians or health care professionals

Data as of August 21, 2018 AS/KAL

Page 8 of 12

This information is considered confidential and proprietary to OptumRx. It is intended for internal use only and should be disseminated only to authorized recipients. The contents of the therapeutic class overviews on this website ("Content") are for informational purposes only. The Content is not intended to be a substitute for professional medical advice, diagnosis, or treatment. Patients should always seek the advice of a physician or other qualified health provider with any questions regarding a medical condition. Clinicians should refer to the full prescribing information and published resources when making medical decisions.

Drug	Available Formulations	Route	Usual Recommended Frequency	Comments
Savella (milnacipran)	Tablet	Oral	Twice daily	Reduce dose in CrCl < 30 mL/min Caution in patients with moderate renal impairment or severe hepatic impairment

Abbreviations: CrCl = creatinine clearance; ESRD = end-stage renal impairment
See the current prescribing information for full details

CONCLUSION

- Included in this review are the neuropathic pain and fibromyalgia agents, duloxetine, gabapentin ER, gabapentin enacarbil ER, lidocaine, pregabalin, pregabalin ER, gabapentin, tapentadol ER, capsaicin, and milnacipran. In general, these agents are FDA-approved for the treatment of diabetic peripheral neuropathic pain, PHN, and/or fibromyalgia.
- Clinical trials support the use of the neuropathic pain and fibromyalgia agents for their FDA-approved indications. Available data demonstrate that neuropathic pain and fibromyalgia agents provide relief from pain; some studies have demonstrated improvement in functional outcomes and quality of life. Direct comparisons among the various agents are rare, and consistent benefit of one agent over another has not been demonstrated.
- According to the available literature, tricyclic antidepressants and duloxetine demonstrate an ability to provide pain relief in patients with painful diabetic neuropathy. While pregabalin and valproate have both demonstrated usefulness in the management of diabetic neuropathy, available literature suggests that the utility of gabapentin is less certain. There is minimal evidence evaluating the use of topical lidocaine for the management of painful diabetic neuropathy. Strong opioids have demonstrated efficacy compared to placebo; however, prescribers may consider this as last line therapy due to concerns regarding long-term safety, including addiction potential and misuse (*Attal et al 2010, Feldman et al 2017[b], Schwartz et al 2011*).
 - Of the neuropathic pain and fibromyalgia agents included in the review, duloxetine, pregabalin, pregabalin ER, and tapentadol ER are approved for the management of diabetic neuropathy.
- For the management of PHN, available literature demonstrates that tricyclic antidepressants, gabapentin, pregabalin, opioids, topical capsaicin, botulinum toxin, and topical lidocaine are more effective compared to placebo (*Bajwa et al 2017*).
 - Of the neuropathic pain and fibromyalgia agents included in this review, gabapentin ER, gabapentin enacarbil ER, lidocaine, pregabalin, pregabalin ER, gabapentin, and capsaicin are approved for the management or relief of pain associated with PHN.
- For the management of fibromyalgia, available literature demonstrates that amitriptyline, cyclobenzaprine, duloxetine, gabapentin, milnacipran, and pregabalin are all appropriate treatment options. The choice of therapy is guided by specific symptoms, comorbidities, and patient preference (*Goldenberg 2016[b]*).
 - Of the neuropathic pain and fibromyalgia agents included in this review, duloxetine and pregabalin are approved for fibromyalgia.

REFERENCES

- American Diabetes Association (ADA). Microvascular Complications and Foot Care: Standards of Medical Care in Diabetes-2018. *Diabetes Care*. 2018;41(Suppl 1):S105-S118. doi: 10.2337/dc18-S010.
- Armstrong DG, Chappell AS, Le TK, et al. Duloxetine for the management of diabetic peripheral neuropathic pain: evaluation of functional outcomes. *Pain Med*. 2007;8(5):410-418.
- Arnold LM, Goldenberg DL, Stanford SB, et al. Gabapentin in the treatment of fibromyalgia: a randomized, double-blind, placebo-controlled, multicenter trial. *Arthritis Rheum*. 2007;56(4):1336-1344.
- Arnold LM, Hudson JI, Wang F, et al. Comparisons of the efficacy and safety of duloxetine for the treatment of fibromyalgia in patients with vs without major depressive disorder. *Clin J Pain*. 2009;25:461-468.
- Arnold LM, Russell IJ, Diri EW, et al. A 14-week, randomized, double-blinded, placebo-controlled monotherapy trial of pregabalin in patients with fibromyalgia. *J Pain*. 2008;9(9):792-805.
- Attal N, Cruccu G, Baron R, et al. EFNS guidelines on the pharmacological treatment of neuropathic pain: 2010 revision. *Eur J Neurol*. 2010;17(9):1113-e88.
- Backonja MM, Canafax DM, Cundy KC. Efficacy of gabapentin enacarbil vs placebo in patients with postherpetic neuralgia and a pharmacokinetic comparison with oral gabapentin. *Pain Medicine*. 2011;12:1098-1108.
- Backonja MM, Wallace MS, et al. NGX-4010, a high-concentration capsaicin patch, for the treatment of postherpetic neuralgia: a randomised, double-blind study. *Lancet Neurol*. 2008;7(12):1106-1112.

- Bajwa ZH. Postherpetic neuralgia. In: Basow DS (Ed). UpToDate [database on the internet]. Waltham (MA): UpToDate; updated February 13, 2017. Available from: <http://www.uptodate.com/contents/postherpetic-neuralgia>. Accessed February 12, 2018.
- Bril V, England J, Franklin GM, et al. Evidence-based guideline: treatment of painful diabetic neuropathy: report of the American Academy of Neurology, the American Association of Neuromuscular and Electrodiagnostic Medicine, and the American Academy of Physical Medicine and Rehabilitation. *Neurology*. 2011;76(20):1758-1765.
- Chou R, Carson S, Chan BK. Gabapentin vs tricyclic antidepressants for diabetic neuropathy and post-herpetic neuralgia: discrepancies between direct and indirect meta-analyses of randomized controlled trials. *J Gen Intern Med*. 2009;24(2):178-188.
- Clauw DJ, Mease P, Palmer RH, et al. Milnacipran for the treatment of fibromyalgia in adults: a 15-week, multicenter, randomized, double-blind, placebo-controlled, multiple-dose clinical trial. *Clin Ther*. 2008;30(11):1988-2004.
- Clauw DJ. Fibromyalgia: an overview. *Am J Med*. 2009;122(12 Suppl):S3-S13.
- Crofford LJ, Rowbotham MC, Mease PJ, et al. Pregabalin for the treatment of fibromyalgia syndrome: results of a randomized, double-blind, placebo-controlled trial. *Arthritis Rheum*. 2005;52(4):1264-1273.
- Crofford LJ. Chapter 396. Fibromyalgia. In: Kasper D, Fauci A, Hauser S, et al. eds. *Harrison's Principles of Internal Medicine*, 19e. New York, NY: McGraw-Hill; 2015. Accessed February 12, 2018.
- Cymbalta [package insert]. Eli Lilly and Company. Indianapolis, IN. December 2017.
- Derry S, Sven-Rice A, Cole P, Tan T, Moore RA. Topical capsaicin (high concentration) for chronic neuropathic pain in adults. *Cochrane Database Syst Rev*. 2017;13(1):CD007393.pub4.
- Drugs@FDA: FDA approved drug products. Food and Drug Administration Web site. <https://www.accessdata.fda.gov/scripts/cder/daf/>. Accessed February 12, 2018.
- Dubinsky RM, Kabbani H, El-Chami, et al; Quality Standards Subcommittee of the American Academy of Neurology. Practice parameter: treatment of postherpetic neuralgia: an evidence-based report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology*. 2004;63:959.
- Dworkin RH, Corbin AE, Young JP Jr, et al. Pregabalin for the treatment of postherpetic neuralgia: a randomized, placebo-controlled trial. *Neurology*. 2003;60:1274-1283.
- Dworkin RH, O'Connor AB, Backonja M, et al. Pharmacologic management of neuropathic pain: evidence-based recommendations. *Pain*. 2007;132(3):237-251.
- Edelsberg JS, Lord C, Oster G. Systematic review and meta-analysis of efficacy, safety, and tolerability data from randomized controlled trials of drugs used to treat postherpetic neuralgia. *Ann Pharmacother*. 2011;45:1483-1490.
- FDA Approved Risk Evaluation and Mitigation Strategies (REMS). [database on the internet]. Extended-Release (ER) and Long-acting (LA) opioid analgesics risk evaluation and mitigation strategy (REMS). Rockville, MD: U.S. Food and Drug Administration. Updated May 2017. Available from: <http://www.accessdata.fda.gov/scripts/cder/remis/index.cfm?event=RemsDetails.page&REMS=17>. Accessed February 12, 2018.
- Feldman EL, McCulloch DK. Treatment of diabetic neuropathy. In: Basow DS (Ed). UpToDate [database on the internet]. Waltham (MA): UpToDate; updated September 20, 2017[b]. Available from: <https://www.uptodate.com/contents/treatment-of-diabetic-neuropathy>. Accessed February 12, 2018.
- Feldman EL. Epidemiology and classification of diabetic neuropathy. In: Basow DS (Ed). UpToDate [database on the internet]. Waltham (MA): UpToDate; updated December 15, 2016[a]. Available from: <https://www.uptodate.com/contents/epidemiology-and-classification-of-diabetic-neuropathy>. Accessed February 12, 2018.
- Fitzcharles MA, Ste-Marie PA, Goldenberg DL, et al; National Fibromyalgia Guideline Advisory Panel. 2012 Canadian Guidelines for the diagnosis and management of fibromyalgia syndrome: executive summary. *Pain Res Manag*. 2013;18(3):119-126.
- Freynhagen R, Strojek K, Griesing T, et al. Efficacy of pregabalin in neuropathic pain evaluated in a 12-week, randomized, double-blind, multicentre, placebo-controlled trial of flexible- and fixed-dose regimens. *Pain*. 2005;115:254-263.
- Galer B, Jensen M, Ma T, et al. The lidocaine patch 5% effectively treats all neuropathic pain qualities: results of a randomized, double-blind, vehicle-controlled, three-week efficacy study with use of the neuropathic pain scale. *Clin J Pain*. 2002;18(5):297-301.
- Galer B, Rowbotham M, Perander J, et al. Topical lidocaine patch relieves postherpetic neuralgia more effectively than a vehicle topical patch: results of an enriched enrollment study. *Pain*. 1999;80:533-538.
- Gilron I, Bailey RN, Tu D, et al. Morphine, gabapentin, or their combination for neuropathic pain. *N Engl J Med*. 2005;352:1324-1334.
- Gralise [package insert]. Depomed, Inc. Newark, CA. September 2015.
- Guan Y, Ding X, Cheng Y, et al. Efficacy of pregabalin for peripheral neuropathic pain: results of an eight-week, flexible-dose, double-blind, placebo-controlled study conducted in China. *Clin Ther*. 2011;33:159-166.
- Handelsman Y, Bloomgarden ZT, Grunberger G, et al. American Association of Clinical Endocrinologists and American College of Endocrinology clinical practice guideline for developing a diabetes mellitus comprehensive care plan- 2015. *Endocrine Practice*. 2015;21(Suppl 1):39-44.
- Hauser W, Bernardy K, Uceyler N, et al. Treatment of fibromyalgia syndrome with antidepressants: a meta-analysis. *JAMA*. 2009[a];301(2):198-209.
- Hauser W, Bernardy K, Uceyler N, et al. Treatment of fibromyalgia syndrome with gabapentin and pregabalin - a meta-analysis of randomized controlled trials. *Pain*. 2009[b];145(1-2):69-81.
- Hauser W, Petzke F, Sommer C. Comparative efficacy and harms of duloxetine, milnacipran, and pregabalin in fibromyalgia syndrome (abstract). *J Pain*. 2010;11(6):505-521.
- Hauser W, Urrutia G, Tort S, et al. Serotonin and noradrenaline reuptake inhibitors (SNRIs) for fibromyalgia syndrome. *Cochrane Database Syst Rev*. 2013;1:CD010292.
- Herndon CM, Strickland JM, Ray JB. Chapter 60. Pain Management. In: DiPiro JT, Talbert RL, Yee GC, Matzke GR, Wells BG, Posey LM, eds. *Pharmacotherapy: A Pathophysiologic Approach*. 10th ed. New York: McGraw-Hill; 2017.
- Horizant [package insert]. Xenoport, Inc. Santa Clara, CA. October 2016.
- Huffman CL, Goldenberg JN, Weintraub J, et al. Efficacy and safety of once-daily controlled-release pregabalin for the treatment of patients with postherpetic neuralgia: a double-blind, enriched enrollment randomized withdrawal, placebo-controlled trial. *Clin J Pain*. 2017;33(7):569-578. doi: 10.1097/AJP.0000000000000445.
- Ifuku M, Iseki M, Hidaka I, et al. Replacement of gabapentin with pregabalin in postherpetic neuralgia therapy. *Pain Medicine*. 2011;12:1112-1126.

- Irving G, Backonja M, Rauck R, et al. NGX-4010, a Capsaicin 8% Dermal Patch, Administered Alone or in Combination With Systemic Neuropathic Pain Medications, Reduces Pain in Patients With Postherpetic Neuralgia. *Clin J Pain*. 2012;28(2):101-107.
- Irving G, Jensen M, Cramer M, et al. Efficacy and tolerability of gastric-retentive gabapentin for the treatment of postherpetic neuralgia: results of a double-blind, randomized, placebo-controlled clinical trial (abstract). *Clin J Pain*. 2009;25(3):185-192.
- Jensen MP, Chiang YK, Wu J. Assessment of pain quality in a clinical trial of gabapentin extended release for postherpetic neuralgia (abstract). *Clin J Pain*. 2009;25(4):286-292.
- Kajdasz DK, Iyengar S, Desai D, et al. Duloxetine for the management of diabetic peripheral neuropathic pain: evidence-based findings from post hoc analysis of three multicenter, randomized, double-blind, placebo-controlled, parallel-group studies. *Clin Ther*. 2007;29:2536-2546.
- Katz N, Gammaitoni A, Davis MW, et al; Lidoderm Patch Study. Lidocaine patch 5% reduces pain intensity and interference with quality of life in patients with postherpetic neuralgia: an effectiveness trial. *Pain Medicine*. 2002;3(4):324-332.
- Lee YH, Song GG. Comparative efficacy and tolerability of duloxetine, pregabalin, and milnacipran for the treatment of fibromyalgia: a Bayesian network meta-analysis of randomized controlled trials. *Rheumatol Int*. 2016;36(5):663-672.
- Lesser H, Sharma U, LaMoreaux L, et al. Pregabalin relieves symptoms of painful diabetic neuropathy. *Neurology*. 2004;63:2104-2110.
- Lidoderm [package insert]. Endo Pharmaceuticals Inc. Malvern, PA. January 2015.
- Lunn MPT, Hughes RAC, Wiffen PJ. Duloxetine for treating painful neuropathy or chronic pain. *Cochrane Database Syst Rev* 2009;(4):CD007115.pub2.
- Lunn MPT, Hughes RAC, Wiffen PJ. Duloxetine for treating painful neuropathy, chronic pain or fibromyalgia. *Cochrane Database Syst Rev* 2014;(1):CD007115.pub3.
- Lyrica [package insert]. Pfizer Inc. New York, NY. May 2018.
- Lyrica CR [package insert]. Pfizer Inc. New York, NY. October 2017.
- Macfarlane GJ, Kronisch C, Dean LE, et al. EULAR revised recommendations for the management of fibromyalgia. *Ann Rheum Dis*. 2017 Feb;76(2):318-328. doi: 10.1136/annrheumdis-2016-209724.
- Mease PJ, Clauw DJ, Gendreau RM, et al. The efficacy and safety of milnacipran for treatment of fibromyalgia: a randomized, double-blind, placebo-controlled trial. *J Rheumatol*. 2009;36:398-409.
- Mease PJ, Russell IJ, Kajdasz DK, et al. Long-term safety, tolerability, and efficacy of duloxetine in the treatment of fibromyalgia. *Semin Arthritis Rheum*. 2010;39:454-464.
- Meier T, Wasner G, Faust M, et al. Efficacy of lidocaine patch 5% in the treatment of focal peripheral neuropathic pain syndromes: a randomized, double-blind, placebo-controlled study. *Pain*. 2003;106:151-158.
- Meng FY, Zhang LC, Liu Y, et al. Efficacy and safety of gabapentin for treatment of postherpetic neuralgia: a meta-analysis of randomized controlled trials. *Minerva Anesthesiol*. 2014;80(5):556-567.
- Micromedex 2.0 [database on the Internet]. Greenwood Village, CO: Truven Health Analytics; 2018. Available from: <http://www.micromedexsolutions.com/home/dispatch>. Accessed February 12, 2018.
- Moon DE, Lee DI, Lee SC, et al. Efficacy and tolerability of pregabalin using a flexible, optimized dose schedule in Korean patients with peripheral neuropathic pain: a 10-week, randomized, double-blind, placebo-controlled, multicenter study. *Clin Ther*. 2010;32:2370-2385.
- Moore RA, Straube S, Wiffen PJ, et al. Pregabalin for acute and chronic pain in adults. *Cochrane Database Syst Rev*. 2009;(3):CD007076.pub2.
- Moore RA, Wiffen PJ, Derry S, et al. Gabapentin for chronic neuropathic pain and fibromyalgia in adults. *Cochrane Database Syst Rev* 2014;(3):CD007938.pub3.
- Neurontin [package insert]. Pfizer Inc. New York, NY. October 2017.
- Nucynta ER [package insert]. Janssen Pharmaceuticals, Inc. Titusville, NJ. June 2017.
- Ogawa S, Suzuki M, Arakawa A, et al. Long-term efficacy and safety of pregabalin in patients with postherpetic neuralgia: results of a 52-week, open-label, flexible-dose study (abstract). *Masui*. 2010;59(8):961-970.
- Orange Book: Approved drug products with therapeutic equivalence evaluations. Food and Drug Administration Web site. <https://www.accessdata.fda.gov/scripts/cder/ob/default.cfm>. Accessed February 12, 2018.
- Parsons B, Li C. The efficacy of pregabalin in patients with moderate and severe pain due to diabetic peripheral neuropathy. *Curr Med Res Opin*. 2016;32(5):929-937.
- Pop-Busui R, Coulton A JM, Feldman EL, Bril V, Freeman R, Malik, R, et al. Diabetic neuropathy: a position statement by the American Diabetes Association. *Diabetes Care*. 2017;40(1):136-154.
- Quilici S, Chancellor J, Lothgren M, et al. Meta-analysis of duloxetine vs pregabalin and gabapentin in the treatment of diabetic peripheral neuropathic pain. *BMC Neurol*. 2009;9:6-19.
- Qutenza [package insert]. Acorda Therapeutics, Inc. Ardsley, NY. August 2013.
- Raskin J, Smith TR, Wong K, et al. Duloxetine vs routine care in the long-term management of diabetic peripheral neuropathic pain. *J Palliative Med*. 2006;9(1):29-40.
- Rice ASC, Maton S; Postherpetic Neuralgia Study Group. Gabapentin in postherpetic neuralgia: a randomized, double blind, placebo controlled study. *Pain*. 2001;94:215-24.
- Rosenstock J, Tuchman M, LaMoreaux L, et al. Pregabalin for the treatment of painful diabetic peripheral neuropathy: a double-blind, placebo-controlled trial. *Pain*. 2004;110:628-638.
- Roth T, van Seventer R, Murphy TK. The effect of pregabalin on pain-related sleep interference in diabetic peripheral neuropathy or postherpetic neuralgia: a review of nine clinical trials. *Clin Med Res & Opin*. 2010;26(10):2411-2419.
- Rowbotham M, Harden N, Stacey B, et al; Gabapentin Postherpetic Neuralgia Study Group. Gabapentin for the treatment of postherpetic neuralgia. A randomized controlled trial. *JAMA*. 1998;280:1837-1842.
- Russell IJ, Mease PJ, Smith TR, et al. Efficacy and safety of duloxetine for treatment of fibromyalgia in patients with or without major depressive disorder: Results from a six-month, randomized, double-blind, placebo-controlled, fixed-dose trial. *Pain*. 2008;136:432-444.
- Sabatowski R, Galvez R, Cherry DA, et al. Pregabalin reduces pain and improves sleep and mood disturbances in patients with post-herpetic neuralgia: results of a randomized, placebo-controlled clinical trial. *Pain*. 2004;109:26-35.
- Savella [package insert]. Forest Pharmaceuticals, Inc. New York, NY. December 2017.

- Schwartz S, Etropolski M, Shapiro DY, et al. Safety and efficacy of tapentadol ER in patients with painful diabetic peripheral neuropathy: results of a randomized-withdrawal, placebo-controlled trial. *Curr Med Res Opin.* 2011;27(1):151-162
- Semel D, Murphy TK, Zlateva G, et al. Evaluation of the safety and efficacy of pregabalin in older patients with neuropathic pain: results from a pooled analysis of 11 clinical studies. *BMC Family Practice.* 2010;11:85.
- Sharma U, Griesing T, Emir B, et al. Time to onset of neuropathic pain reduction: a retrospective analysis of data from nine controlled trials of pregabalin for painful diabetic peripheral neuropathy and postherpetic neuralgia. *Am J Ther.* 2010;17:577-585.
- Siddall PJ, Cousins MJ, Otte A, et al. Pregabalin in central neuropathic pain associated with spinal cord injury: a placebo-controlled trial. *Neurology.* 2006 Nov 28;67(10):1792-1800.
- Skvarc NK, Kamenik M. Effects of pregabalin on acute herpetic pain and postherpetic neuralgia incidence. *Wien Klin Wochenschr.* 2010;122(Suppl 2):49-53.
- Tanenberg RJ, Clemow DB, Giaconia JM, et al. Duloxetine compared with pregabalin for diabetic peripheral neuropathic pain management in patients with suboptimal pain response to gabapentin and treated with or without antidepressants: a post hoc analysis. *Pain Pract.* 2014;14(7):640-648.
- Tanenberg RJ, Irving GA, Risser RC, et al. Duloxetine, pregabalin, and duloxetine plus gabapentin for diabetic peripheral neuropathic pain management in patients with inadequate pain response to gabapentin: an open-label, randomized, noninferiority comparison. *Mayo Clin Proc.* 2011;86(7):615-24.
- Tesfaye S, Wilhelm S, Lledo A, et al. Duloxetine and pregabalin: High-dose monotherapy or their combination? The "COMBO-DN study" – a multinational, randomized, double-blind, parallel-group study in patients with diabetic peripheral neuropathic pain. *Pain.* Dec 2013;154(12):2616-2625.
- Vinik AL, Perrot S, Vinik EJ, et al. Capsaicin 8% patch repeat treatment plus standard of care (SOC) versus SOC alone in painful diabetic peripheral neuropathy: a randomised, 52-week, open-label, safety study. *BMC Neurol.* 2016;16(1):1-14.
- Vitton O, Gendreau M, Gendreau J, et al. A double-blind placebo-controlled trial of milnacipran in the treatment of fibromyalgia. *Hum Psychopharmacol Clin Exp.* 2004;19:S27-S35.
- Vranken JH, Kijkgraaf MG, Kruis MR, et al. Pregabalin in patients with central neuropathic pain: a randomized, double-blind, placebo-controlled trial of a flexible-dose regimen. *Pain.* 2008;136(1-2):150-157.
- Wallace MS, Irving G, Crowles VE. Gabapentin extended-release tablets for the treatment of patients with postherpetic neuralgia: a randomized, double-blind, placebo-controlled, multicentre study (abstract). *Clin Drug Investig.* 2010;30(11):765-776.
- Wernicke J, Lledo A, Raskin J, et al. An evaluation of the cardiovascular safety profile of duloxetine. *Drug Safety.* 2007[a];30(5):437-455.
- Wernicke J, Wang F, Pritchett YL, et al. An open-label 52-week clinical extension comparing duloxetine with routine care in patients with diabetic peripheral neuropathic pain. *Pain Medicine.* 2007[b];8(6):503-513.
- Williams DA, Schilling S. Advances in the assessment of fibromyalgia. *Rheum Dis Clin North Am.* 2009;35(2):339-57.
- Xochilcal-Morales M, Castro EM, Guajardo-Rosas J, et al. A prospective, open-label, multicentre study of pregabalin in the treatment of neuropathic pain in Latin America. *Int J Clin Pract.* 2010;64(9):1301-1309.
- Yan G, Guang N, Wei-ping J, et al. Duloxetine vs placebo in the treatment of patients with diabetic neuropathic pain in China. *Chin Med J.* 2010;123(22):3184-3192.
- Zhang L, Rainka M, Freeman R, et al. A randomized, double-blind, placebo-controlled trial to assess the efficacy and safety of gabapentin enacarbil in subjects with neuropathic pain associated with postherpetic neuralgia (PXN110748). *J Pain.* 2013;14(6):590-603.
- ZTlido [package insert]. Scilex Pharmaceuticals Inc. San Diego, CA. February 2018.

Publication Date: August 22, 2018